

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
1 July 2004 (01.07.2004)

PCT

(10) International Publication Number
WO 2004/054615 A1

- (51) International Patent Classification⁷: **A61K 39/395**, 47/48, 51/10 // A61M 1/36 (74) Agent: AWAPATENT AB; Box 5117, S-200 71 Malmö (SE).
- (21) International Application Number: PCT/SE2003/001949
- (22) International Filing Date: 12 December 2003 (12.12.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0203731-5 13 December 2002 (13.12.2002) SE
60/433,012 13 December 2002 (13.12.2002) US
- (71) Applicant (for all designated States except US): MITRA MEDICAL TECHNOLOGY AB [SE/SE]; Ideon Research Park, S-223 70 Lund (SE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): SANDBERG, Bengt [SE/SE]; Blomstergården 18, S-245 62 Hjärrup (SE). NILSSON, Rune [SE/SE]; Utsättaregården 139, S-226 47 Lund (SE).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, EG, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTILYMPHOMA TARGETING AGENTS WITH EFFECTOR AND AFFINITY FUNCTIONS LINKED BY A TRIFUNCTIONAL REAGENT



(57) Abstract: A medical agent comprising a reagent conjugated to an anti-lymphoma antibody is disclosed, as well as a kit containing said medical agent, use of said medical agent, and a method for treatment of lymphoma. The reagent may comprise an effector, e.g. an antitumor agent or a diagnostic marker, and an affinity ligand enabling extracorporeal clearance of the agent. The three components are bound by a trifunctional linker.

WO 2004/054615 A1

This Page Blank (uspto)

Antilymphoma targeting agents with effector and affinity functions linked by a trifunctional reagent.

Technical Field of the Invention

The present invention relates to a medical agent comprising a reagent conjugated to an anti-lymphoma antibody, a kit for treating or diagnosing lymphoma, use
5 of said medical agent, and a method for treatment of lymphoma.

Background Art

Lymphomas are malignant cell infiltrations of the lymphatic system. The lymphatic system includes the nodes
10 which are located in the neck, armpit, and groin. These nodes are only part of the lymphatic system, as they are connected to each other and to the spleen, thymus, and parts of the tonsils, stomach, and small intestine by a network of vessels. The vessels carry a fluid called
15 lymph which contains lymphocytes. Once a malignancy begins in one part of the lymphatic system, it often spreads throughout the rest of the lymphatic system before it is detected.

There are precise, internationally agreed criteria
20 to define the stage of disease for each type of cancer. For lymphomas this means mapping out how many lymph nodes are affected. It also means finding out if the lymphoma has spread outside the lymphatic system to other organs. Stage I: Cancer limited to one group of lymph nodes or a
25 single organ or site outside the lymphatic system Stage II: Cancer in two or more groups of lymph nodes all on the same side of the diaphragm Stage III: Cancer on both sides of the diaphragm but not outside the lymphatic system
30 Stage IV: Widespread cancer outside the lymphatic system

Lymphomas are divided into many sub-groups according to cell types. Generally, they are classified as non-Hodgkin's and Hodgkin's. Currently, Hodgkin's lymphoma is more curable than non-Hodgkin's. Non-Hodgkin's lymphomas

are derived from both B-cells and T-cells origins, where 90% of all cases are B-cell derived and the remaining 10% are of T-cell derivation.

5 The treatment for all types of lymphoma depends on the type, stage, and grade of disease. The stages and grades are outlined below.

Stages:

- I: cancer site, no bone marrow involvement
- 10 II: two sites, both either above or below the diaphragm; no bone marrow involvement
- III: sites above and below the diaphragm; no bone marrow involvement
- IV: bone marrow is affected or the cancer cells have spread outside the lymphatic system

15

Grades:

- high: usually found in B-cell and T-cell types
- intermediate: usually found in B-cell and T-cell types
- 20 low: predominantly found in B-cell types

Lymphomas are usually treated by a combination of chemotherapy, radiation, surgery, and/or bone marrow transplants. The cure rate varies greatly depending on
25 the type of lymphoma and the progression of the disease.

Because lymph tissue is found in many parts of the body, non-Hodgkin's lymphoma can start in almost any part of the body. The cancer can spread to almost any organ or tissue in the body, including the liver, bone marrow,
30 spleen, and nose.

Based on the histology, non-Hodgkin's lymphomas are divided into two groups: indolent lymphomas, which grow more slowly and have fewer symptoms, and aggressive lymphomas, which grow more quickly.

35 Lymphomas include follicular small cleaved cell lymphoma, adult diffuse mixed cell lymphoma, follicular mixed cell lymphoma, adult diffuse large cell lymphoma,

follicular large cell lymphoma, adult immunoblastic large cell lymphoma, adult diffuse small cleaved cell lymphoma, adult lymphoblastic lymphoma, small lymphocytic (marginal zone) adult small non-cleaved cell lymphoma.

5 Other types of indolent non-Hodgkin's lymphoma/-leukemia are lymphoplasmacytoid lymphoma, monocytoid B-cell lymphoma, mucosa-associated lymphoid tissue (MALT) lymphoma, splenic marginal zone lymphoma, hairy cell leukemia, and cutaneous T-cell lymphoma (Mycosis
10 fungoides/Sezary syndrome).

Other types of aggressive non-Hodgkin's lymphoma are anaplastic large-cell lymphoma, adult T-cell lymphoma/-leukemia, mantle cell lymphoma, intravascular lymphomatosis, angioimmunoblastic T-cell lymphoma, angiocentric
15 lymphoma, intestinal T-cell lymphoma, primary mediastinal B-cell lymphoma, peripheral T-cell lymphoma, lymphoblastic lymphoma, post-transplantation lymphoproliferative disorder, true histiocytic lymphoma, primary central nervous system lymphoma, and primary effusion lymphoma.
20 Aggressive lymphomas are also seen more frequently in patients who are HIV-positive (AIDS-related lymphoma).

Recurrent adult non-Hodgkin's lymphoma may come back in the lymph system or in other parts of the body.

Indolent lymphoma may come back as aggressive lymphoma. Aggressive lymphoma may come back as indolent
25 lymphoma.

Non-Hodgkin's lymphomas (NHLs) are the fifth leading cause of cancer morbidity and mortality (Wingo P, Tong T, Bolden S. Cancer statistics, 1995. CA Cancer J. clin
30 1995; 45, 8-30 & Parker SL, Tong T, Bolden S., Wingo PA. Cancer statistics, 1996. CA Cancer J. Clin 1996;46:5-27). Over the past two decades, the prevalence in the USA of these lymphomas has increased rapidly. Five years ago more than 52,000 new diagnosis were made, and 23.000
35 deaths were attributed to NHLs and with an incidence increasing at a rate of 7% per year (Parker SL, Tong T, Bolden S., Wingo PA. Cancer statistics, 1996. CA Cancer

J. Clin 1996;46:5-27). This represents an increase of 150% in the population-adjusted new cases of NHLs over the past 50 years.

The overwhelming majority of patients (about 80%) constitute patients with NHLs of B-cell origin (Harris N.L., Jaffe E.S., Stein H. et.al.. Lymphoma classification proposal: clarification [letter]. Blood 1995; 85: 857-860). Despite the use of various combined chemotherapeutic regimens for advanced-stage intermediate- and high grade lymphomas, roughly half of patients treated do not have a complete remission or finally have a relapse after remission. The situation has not improved noticeably in almost two decades (Gordon LI, Harrington D, Andersen J, et.al. N. Engl. J. Med. 1992;327:1342-9 & Fisher RI, Gaynor ER, Dahlberg S, et.al. N. Engl. J. Med. 1993; 328: 1002-6).

Treatment with standard-dose salvage chemotherapy rarely results in durable remissions and often has serious toxicity. Although the use of high-dose chemotherapy with bone marrow transplantation has shown to be promising, not all patients derive long-term benefits from this type of treatment (Armitage JO. Blood 1989;73:1749-58). A curative treatment for patients with advanced low-grade lymphoma still remains to be clearly established (DeVite VT Jr., Jaffe ES, Mauch P, Longo DL. Lymphocytic lymphomas. In: DeVita VT Jr., Hellman S., Rosenberg SA. Eds. Cancer: principles and practice of oncology. 3rd ed. Vol.2. Philadelphia: J.B. Lippincott, 1989;1741-98). Treatment with anthracycline-based chemotherapy regimes results in complete remission in 50-90 percent of patients with intermediate and high-grade non-Hodgkin's lymphoma and long-term disease-free survival in 30-60 percent.

Unfortunately, few patients with low-grade lymphoma or relapses of any type of lymphoma can be cured with conventional approaches (Armitage JO. N.Engl.J.Med. 1993; 328:1023-30). High-dose chemoradiotherapy with bone

marrow transplantation cures 10-50% of patients with lymphoma in relapse, but 40-80% relapse again and 5-20% die of complications related to transplantation (Appelbaum FR, Sullivan KM, Buckner CD et.al. J.Clin. Oncol. 1987;5:1340-7 & Freedman AS, Takvorian T, Anderson KC et.al. J.Clin.Oncol. 1990;8:784-91). The use of large doses of chemoradiotherapy has not been feasible because of unacceptable morbidity and mortality (Bearman SI, Appelbaum FR, Bruchner CD. et.al. J.Clin.Oncol. 1988; 6:1562-8).

Tissue or organ specific localisation of a medical agent is a very important factor in its effective application. Lack of specific tissue localisation is of particular importance in the treatment with cytotoxic agents, where the desired effect is to kill certain types of cells, such as in the treatment of cancer.

The treatment of cancer with agents specific for the tumour cell without harming the host has long been a goal of oncology. The development of monoclonal antibodies provided hope that tumour-targeted therapy would one day play a role in the treatment of cancer. Indeed, promising results have been presented in several areas; however, most of the treatment modalities have often proved technically difficult, produced disappointing efficacy, and were often not broadly applicable to patients with a given malignancy.

The treatment of patients with lymphoma is an exception. Patients with advanced stage or relapsed low-grade non-Hodgkin's lymphoma (NHL) are not curable using conventional approaches and are usually treated with combination chemotherapy regimens of increasing intensity as needed to reduce disease and palliate symptoms. Recent attempts utilising supralethal chemotherapy combined with radiotherapy followed by bone marrow transplantation have resulted in an approximately 20% long term disease-free survival rate (F. Applebaum et al, J. Clin.Oncol. 5:1340, 1987). However, most patients treated in this manner die

of lymphoma or treatment complications. Therefore, new strategies for the treatment of non-Hodgkin's lymphomas are needed. These strategies should be aiming at the maximisation of therapeutic effect coupled with the minimization of toxicity.

One approach involves the use of monoclonal antibodies that recognise tumour-associated antigens as a means of targeting drugs or radioisotopes to tumour cells. This approach is particularly attractive in the case of NHL as the lymphoma tumour cells display a variety of tumour-restricted antigens on their cell surfaces that would be available for targeting (A.J. McMichael, Leukocyte Typing III, pp 302-363 and 432-469, Oxford University Press, Oxford, England, 1987). The rationale for utilising such an approach is further supported by the observation that monoclonal antibodies by themselves can exhibit anti-tumour effects in vivo. Of all the malignancies that have been treated with monoclonal antibodies to date, the lymphomas have yielded the most dramatic results. Significant tumour regressions have been reported in patients treated with monoclonal anti-idiotypic antibodies (R.A Miller et al, New Eng. J. Med. 306:517, 1982; T.C. Meeker et al, Blood 65:1349, 1985). Most of the tumour responses, however, have been incomplete and of relatively short duration. The practical problem of generating anti-idiotypic antibodies restricts the utility of such an approach (T. Meeker et al, New Eng. J. Med 312:1658, 1985).

Recently, a number of monoclonal antibodies have been developed which recognise antigenic sites on both malignant and normal human B cells. These pan B-cell antibodies have been useful in classifying lymphomas and in defining the ontogeny and biology of normal B cells. Because of the limited efficacy of unmodified antibodies in general, recent attention has focused on the use of antibodies conjugated to cytotoxic agents. Among the cytotoxic agents that might be considered, radioisotopes

are especially attractive, as lymphomas are especially sensitive to the effects of radiation. Moreover, such radiolabelled antibodies may be of considerable utility in terms of diagnostic imaging of tumour involved sites.

5 Most of these cytotoxic anti-lymphoma antibodies are directed towards CD20.

CD20 is an antigen that is a 35 kilodaltons, non-glycosylated phosphoprotein found on the surface of greater than 90% of B cells from peripheral blood or lymphoid
10 organs. The antigen is expressed on the surface of virtually all resting B cells maintained in culture, but is lost by approximately one-third of the population upon activation of the cells by protein A or exposure to Epstein-Barr virus. This result has been interpreted to
15 mean that CD20 is lost during terminal differentiation of B cells (L. M. Nadler, Lymphocyte typing II, vol 2 pp 3-37 and 65 Appendix, E. L. Renling et al eds Springer Verlag, 1986).

A number of other antigens like the CD19 are also
20 expressed on the surface of cells of the B lineage. However, contrary to the CD20, antibodies binding to the CD19 are rapidly internalised. Other antibodies identified as binding to these types of cells are: the B2 binding to the CD21 antigen; B3 binding to the CD22 antigen
25 and the J5 binding to the CD 10 antigen. The pan-B-cell antibody MB-1 is also of interest and has been shown to bind to CD37.

Naked antibodies directed against CD20 have shown to have efficiency. One registered naked antibody,
30 Rituximab, is a chimeric mouse/human anti-CD20 antibody that has shown efficiency in the treatment of indolent lymphoma, especially follicular lymphoma. The overall response rate for patients with indolent lymphoma is 50% and the complete response rate is 10% (McLaughlin P et
35 al, J. Clin. Oncol. 16: 2825-2833, 1998.). Time to progression has been reported to be 13 months. Rituximab does also produce objective remissions in aggressive

lymphoma albeit with a lower response rate. Nonetheless virtually all patients treated with Rituximab as a single agent will finally relapse.

Systemic radiotherapy is an established form of treatment. The use of radioiodine in the treatment of disseminated cancer of the thyroid is often the mainstay of therapy. Radioimmunotherapy (RIT) is another form of systemic radiotherapy where the radionuclide is targeted by an antibody to a tumour cell. RIT is in some cases a combined modality between radiotherapy and immunotherapy, since the antibody itself may exert an anti-tumour effect. The use of RIT is still experimental, but several encouraging studies have been published. In treatment of B-cell lymphoma several groups have reported long term remissions following RIT. Most investigators have used ^{131}I or ^{90}Y labelled mouse antibodies directed to the CD20 antigen (Kaminski, M. S. et al, J. Clin. Oncol., 14:1974 -1981, 1996, Knox, S. J et al, Clin. Cancer Res., 2: 457-470, 1996.).

Therapeutic application of chimeric and radiolabelled antibodies for treatment of B cell lymphoma is described by Anderson, D.R. et.al. in EP 0 752 248 B1; EP 669 836 B1 and US 5,843,439; 5,776,456; 5,736,137. Methods for the treatment of lymphoma by administration of a B cell-specific antibody are described in Kaminski, M.S. et.al. US 5,595,721; 6,015,542; 5,843,398; 6,090,365 and by Goldenberg, D.M. et.al. in US 6,183,744 B1. Other patents and patent applications related to the subject matter are US 6,399,061 B1, EP 1 005 870 A2, WO 98/42378, WO 99/57981, WO 00/09160, WO 00/27428, WO 00/27433, WO 01/34194 A1, WO 01/10462 A1, WO 01/10460 A1, WO 00/67795, WO 00/52473.

Rituximab is a chimeric mouse/human antibody that has been engineered from its mouse parental antibody, ibritumomab. When ibritumomab is labelled with ^{90}Y , it is entitled ZevalinTM. Wiseman et.al. Critical reviews in Oncology/Hematology 39 (2001), 181-194, have reported

that Zevalin™ may be administered safely without prior dosimetry at an activity of 15 MBq/kg to patients with a platelet count of $> 149 \times 10^9$ /L. For patients with platelet counts of $100-149 \times 10^9$, an activity of 11.1 MBq/ kg is well tolerated. A prospective randomised trial of Zevalin in patients with relapsed or refractory indolent or transformed lymphoma compared to a standard course of Rituximab has been reported. Among 143 patients studied, an overall response of 80% was found for the Zevalin group vs 56% in the group who received unlabelled Rituximab ($P=0.002$) and with 30% complete remission with Zevalin vs 16% CR for Rituximab ($P=0.04$). Zevalin has also been evaluated in patients with follicular lymphoma refractory to Rituximab. The response duration was significantly longer (8.4+ vs 4 months) for Zevalin as compared with prior Rituximab ($P=0.008$).

Tositumomab is a murine IgG_{2a} lambda monoclonal antibody directed against the CD20 antigen. I¹³¹-tositumomab (Bexxar®) is a radio-iodinated derivative of tositumomab that has been covalently linked to Iodine-131. Iodine-131 decays with beta and gamma emissions with a physical half-life of 8.04 days. Possible mechanisms of action of the I¹³¹-tositumomab therapeutic regimen include induction apoptosis, complement dependent cytotoxicity (CDC) (and antibody-dependent cellular cytotoxicity (ADCC) mediated by the antibody) (Cardarelli PM et. al. Cancer Immunol Immunother. 2002 Mar; 51(1): 15-24; Stashenko P, et. al. J Immunol 1980; 125:1678-85). Additionally, cell death is associated with ionizing radiation from the radioisotope.

The therapeutic regimen is administered in two discrete steps: the dosimetric and the therapeutic step. Each step consists of a sequential infusion of tositumomab, followed by I¹³¹-tositumomab.

The maximum dose of the I¹³¹-tositumomab therapeutic regimen that was administered in clinical trials was 88 cGy. Three patients were treated with a total body dose

of 85 cGy of Iodine I¹³¹-tositumomab in a dose escalation study. Two of the 3 patients developed grade 4 toxicity of 5 weeks duration with subsequent recovery. In addition, accidental overdose of the therapeutic regimen
5 occurred in one patient at total body doses of 88 cGy.

Normal organ toxicity limits the amount of activity that can be safely administered to patients and thereby the absorbed dose to tumour. The first dose-limiting organ is the bone marrow. Localised B-cell lymphoma may
10 be cured by external beam radiotherapy with a dose of 30 to 44 Gy. The dose that may be achieved with conventional radioimmunotherapy without the use of stem cell support is substantially lower. Wiseman et al has reported a median dose of 15 Gy in B-cell lymphoma in a phase III
15 trial (Wiseman G et al., Critical reviews in Oncology/-Hematology 39 (2001) 181-194). The response rate was 80% objective response and 34% complete response. The Seattle group using stem cell support has reported the highest remission rate 80% complete remissions (Liu
20 Steven Y. et al., J. Clin. Oncol.16(10): 3270-3278, 1998). They estimated tumour sites to achieve 27 to 92 Gy.

The non-haematological dose-limiting toxicity was reversible pulmonary insufficiency, which occurred at
25 doses \geq 27 Gy to the lungs. Although the studies are not quite comparable, they indicate a dose effect relationship in RIT. If there is a dose relationship, it may be possible to increase efficacy if a higher dose to the tumour can be delivered. This may be most clinically relevant, since complete remission following RIT has been
30 associated with longer duration of remission (Wahl et al., J.Nucl. Med.39:21S-26S, 1998.).

An obstacle to this is the radio sensitivity of the bone marrow. A higher absorbed dose to the bone marrow
35 may cause myeloablation. Thus, the dose necessary to reach a more effective therapy is hampered by the accumulation of radioactivity in the blood circulation,

leading to toxicity of normal organs, such as bone marrow. Various means for clearing blood from cytotoxic targeting biomolecules (e.g. therapeutic or diagnostic monoclonal antibodies) after intravenous administration have been reported (See review article by Schriber G.J. and Kerr D. E., Current Medical Chemistry 2:616-629, (1995)).

In the so-called avidin chase modality, avidin or streptavidin is administered systemically after administration of the therapeutic or diagnostic antibody to which biotin has been attached, at a time when a sufficient amount of the antibody has been accumulated in the tumour. Avidin or streptavidin will associate with the antibodies and the so formed immunocomplex will clear from the blood circulation via the reticuloendothelial system (RES) and be cleared from the patient via the liver. These procedures will improve the clearance of biotinylated cytotoxic antibodies. An alternative approach to the same end is the use of anti-idiotypic antibodies. However, all these methods rely on the liver or kidney for blood clearance and thereby expose either or both of these vital organs as well as the urinary bladder to a high dose of cytotoxicity.

Another major drawback of the methods is the immunogenicity of these agents, particularly the streptavidin, which prevent repetitive treatments once the immune response has been developed. Extracorporeal techniques for blood clearance are widely used in kidney dialysis, where toxic materials build up in the blood due to the lack of kidney function. Other medical applications, whereby an extracorporeal apparatus can be used, include: removal of radioactive materials; removal of toxic levels of metals, removal of toxins produced from bacteria or viruses; removal of toxic levels of drugs, and removal of whole cells (e.g. cancerous cells, specific haematopoietic cells - e.g. B, T, or NK cells) or removal of bacteria and viruses.

Various methods have been proposed to rapidly clear radiolabelled antibodies from blood circulation after the tumour has accumulated a sufficient quantity of immunoconjugate to obtain a diagnosis or therapy. Some of the methods employed involve enhancement of the body's own clearing mechanism through the formation of immune complexes. Enhanced blood clearance of radiolabelled antibodies can be obtained by using molecules that bind to the therapeutic antibody, such as other monoclonal antibodies directed towards the therapeutic antibody (Klibanov et al, J. Nucl. Med 29:1951-1956 (1988); Marshall et al, Br. J. Cancer 69: 502-507 (1994); Sharkey et al, Bioconjugate Chem. 8:595-604, (1997), avidin/streptavidin (Sinitsyn et al J. Nucl. Med. 30:66-69 (1989), Marshall et al Br. J. Cancer 71:18-24 (1995), or glycosyl containing compounds which are removed by receptors on liver cells (Ashwell and Morell Adv. Enzymol. 41:99-128 (1974). Still other methods involve removing the circulating immunoconjugates through extracorporeal methods (See review article by Schreiber G.J. and Kerr D.E., Current Medical Chemistry 2:616-629 (1995)).

The extracorporeal techniques used to clear a medical agent from blood circulation are particularly attractive because the toxic material is rapidly removed from the body.

Applications of these methods in the context of immunotherapy have been previously described (Henry Chemical Abstract 18:565 (1991); Hofheinz D. et al Proc. Am. Assoc. Cancer Res. 28:391 (1987); Lear J. K. et al Antibody Immunoconj. Radiopharm. 4:509 (1991); Dienhart D. G. et al Antibody Immunoconj. Radiopharm. 7:225 (1991); DeNardo S.J. et al J. Nucl. Med 33:862-863 (1992); DeNardo G.L. et al J. Nucl. Med 34:1020-1027 (1993); DeNardo G. L. J. Nucl. Med 33:863-864 (1992); and US patent No. 5,474,772 (Method of treatment with medical agents).

To make the blood clearance more effective and to enable processing of whole blood, rather than blood plasma which the above methods refer to, the medical agents (e.g. tumour specific monoclonal antibody carrying cell killing agents or radio nuclides for tumour localization) have been biotinylated and cleared by an avidin-based adsorbent on a column matrix. A number of publications provide data showing that this technique is both efficient and practical for the clearance of biotinylated and radionuclide labelled tumour specific antibodies (Norrgren K. et al, Antibody Immunoconj. Radiopharm. 4:54 (1991), Norrgren K. et al J. Nucl. Med 34:448-454 (1993); Garkavij M. et al Acta Oncologica 53:309-312 (1996); Garkavij M. et al, J. Nucl. Med. 38:895-901 (1997)).

These techniques are also described in EP 0 567 514 and US 6,251,394. The device Mitradep[®], developed and manufactured by Mitra Medical Technology AB, Lund, Sweden, is based on this technology. By using the avidin-coated filter in conjunction with biotin labelled therapeutic antibodies, the blood clearance technique can be applied equally well to chimeric or fully humanised antibodies. Experimental data reveal that during a three-hour adsorption procedure, more than 90% of the circulating biotinylated antibodies can be removed by the Mitradep[®] system (Clinical Investigator's Brochure - Mitradep[®]).

In order to be adsorbed to the extracorporeal filter, the monoclonal antibodies carrying the cytotoxic agent (e.g. radionuclide) need to be biotinylated (biotin binds irreversible to the avidin in the filter) prior to administration to the patient. The number of biotinyl moieties per IgG molecule is in the range of 3-6, typically 4.

A further development of this method with simultaneous labelling of biotin and radionuclides is described in a patent application by S. Wilbur and B.E.B.

Sandberg PCT/ SE98/01345, disclosing a trifunctional reagent for the conjugation to a biomolecule.

The latter method has a number of advantages over the consecutive labeling of radio nuclides and biotinylation and is particularly attractive in cases where the naked (non-chelated) antibody is supplied to the hospital, since both the chelating group and the biotinyl groups have to be conjugated to the antibody in addition to the radiolabelling step.

However, in most cases the same type of functions (ϵ -amino groups) on the antibodies are utilized for coupling of the chelating groups and the biotinyl groups, leading to a competition of the most accessible sites.

Chelation and/or biotinylation of an antibody results in a heterogenous preparation. If for example a chelated antibody is determined to have 3 chelates per antibody, the preparation contains a mixture of antibodies with 1 chelate/antibody to 7 chelates/antibody. As the chelate and biotin are linked to the same moieties on the antibody, antibodies with a higher number of chelates might have a lower number of biotin. It might also results in antibodies with a high number of chelates having no biotin at all.

This means that, statistically, a population of the antibodies carrying radionuclide but not biotin will circulate in the blood, and those antibodies will not be removed by the Mitradep® filter.

To facilitate the labelling of the naked therapeutic or diagnostic antibody and to ensure that the ratio of biotin and the radiolabel is one to one, Mitra Medical Technology AB, Lund, Sweden has developed a series of novel water-soluble structures (Tag-reagent; MitraTag™) containing the two types of functions, thereby enabling simultaneous and site specific conjugation of chelating groups (for radiolabelling) and the biotin groups.

The Tag-reagent labelled with the chelating group DOTA is called MitraTag-1033.

The present invention encompasses a medical agent comprising a reagent conjugated to an anti-lymphoma antibody, and various methods for the treatment of lymphatic cancer, i.e. lymphoma, and NHL in particular.

5 Summary of the Invention

The object of the present invention is to solve the above discussed problem in connection with treatment of certain lymphoma diseases. This object is achieved by the present invention as specified below.

10 The present invention relates in one aspect to a medical agent comprising a reagent conjugated to an anti-lymphoma antibody or a variant thereof, wherein the reagent is a single molecule with at least three functional parts b) - d) wherein,

- 15 a) a trifunctional cross-linking moiety is coupled to
b) an affinity ligand via a linker 1, to
c) an effector agent via a covalent bond, optionally via a linker 2, and to
d) a biomolecule reactive moiety, optionally via a
20 linker 3, wherein said biomolecule reactive moiety is an anti-lymphoma antibody reactive moiety being capable of forming a bond with the anti-lymphoma antibody or a variant thereof, thereby forming a conjugate, and wherein the anti-lymphoma antibody or
25 variants thereof is/are interacting with one or more different CD antigen(s) present on the surface of lymphoma tumour cells.

In another aspect, the present invention relates to a composition comprising said medical agent.

30 In a further aspect, the present invention relates to a kit for extracorporeal elimination or at least reduction of the concentration of a non-tissue-bound therapeutic or diagnostic medical agent as defined above in the plasma or whole blood of a mammalian host, wherein
35 said medical agent previously has been introduced into a mammalian host and kept therein for a certain time in

order to be concentrated to the specific tissues or cells by being attached thereto, said host comprising

- a) the medical agent, and
- b) an extracorporeal device comprising an immobilised
5 receptor onto which the affinity ligand adheres.

In a further aspect, the present invention relates to use of said medical agent for treatment of lymphoma, preferably non-Hodgkin's lymphoma.

In still a further aspect, the present invention
10 relates to a method for treatment of lymphoma, preferably non-Hodgkin's lymphoma, by the administration of the medical agent.

In still a further aspect, the present invention relates to a method for diagnosing lymphoma, preferably
15 non-Hodgkin's lymphoma, by the administration of the medical agent, as well as to a method for combined treatment and diagnosing lymphoma.

Further advantages and objects of the present invention will now be described in more detail with reference
20 to the accompanying drawings.

Brief Description of the Drawings

Fig. 1 shows depletion of 1033-rituximab conjugates during recirculation through a miniaturised Mitradep®.

Fig. 2 shows a flow cytometric assay of binding to
25 the CD20 positive cell line Raji.

Fig. 3 shows binding of 1033-conjugates to a CD20+(SB) and a CD20-(HSB) cell line.

Fig. 4 shows competitive inhibition of ¹²⁵I-labelled rituximab binding to SB cells by cold rituximab and 1033-
30 rituximab conjugates.

Fig. 5 shows whole body clearance of radioactivity in rats injected with ¹¹¹In-1033-rituximab antibody conjugates expressed as percentage \pm std.dev.

Fig. 6 shows blood clearance of ¹¹¹In-1033-rituximab
35 antibody conjugates expressed as % injected dose/gram \pm std.dev.

Fig. 7 shows biodistribution of ^{111}In -1033-rituximab (4.6 1033/IgG) in rats.

Fig. 8 shows HPLC size exclusion separation of blood samples drawn from a rat injected with ^{111}In -1033-rituximab (4.6 1033/IgG).

Description of Preferred Embodiments

With the present invention it is possible to improve the tumour to non-tumour ratio of cytotoxic targeting agents in the treatment of disseminated haematological carcinomas, in particular lymphomas, by reducing the concentration of the cytotoxic medical agent in the blood circulation after administration of a cytotoxic agent and thereby facilitating a higher dosage and hence a more effective treatment regime without exposing the vital organs to higher toxicity. Furthermore, the present invention presents new medicals and the use of these agents in the treatment of lymphatic cancer and NHL, in particular.

In one embodiment, a radiolabelled anti-lymphoma antibody is given in a single dose which is limited to what is regarded as tolerable to the patient without reconstitution of the hematopoietic function, through bone marrow transplantation, or by some other means; "low dose". The dose range will be 10-20 MBq/kg body weight of ^{90}Y -anti-lymphoma antibody, preferably 11-15 MBq/kg and the range for ^{111}In -anti-lymphoma antibody for targeting localisation will be 20-250 MBq, preferable 50-150 MBq. In this embodiment, extracorporeal clearance of non-bound radiolabelled therapeutic or diagnostic antibody is optional.

In another embodiment, a radiolabelled anti-lymphoma antibody is given in a single dose designated to deliver a high amount of radioactivity to the patient. This "high dose method" has to be combined with means of reconstituting the bone marrow or by reducing the radiation effect on bone marrow preferably by the use of the Mitradep® system. For ^{90}Y -anti-lymphoma antibodies, "high

dose" means a single dose exceeding 20 MBq/kg body weight.

In a preferred embodiment, ^{111}In -anti-lymphoma at a dose of 50-150 MBq is combined with a "high dose" (> 20 MBq/ kg body weight) of ^{90}Y -anti-lymphoma antibody, either
5 given in sequence at intervals of 6-8 days or given simultaneously.

The following embodiments of the invention also serve to explain the details of the invention.

10 Lymphomas are tumours originating from lymphocytes. The normal counterparts of lymphomas, i.e. the normal lymphocytes, arise from pluripotent stem cells in the bone marrow and differentiate to fully mature lymphocytes. During their differentiation they express different cell surface antigens (CD-antigens) some of which
15 are lineage and/or stage specific. Lymphomas can arise from lymphocytes in various differentiation stages and often present the CD-antigens expressed at this stage. These CD-antigens cannot be used only for diagnostic
20 purposes but also as targets for different kinds of antibody therapy.

The study of human leukocyte antigens, predominantly by monoclonal antibody techniques, is a rapidly changing field of basic research and clinical investigation.
25 Leukocyte surface molecules defined by antibodies have been assigned cluster differentiation (CD) numbers (CD-antigens) in a series of international workshops (Paris, 1982; Boston, 1984; Oxford, 1986; Vienna, 1989, Osaka, 1996). The CD classification of these antigens has
30 become the standard form in published literature and provides a basis for standardization of clinical reporting. The current CD classification is presented in the form of a list, with a brief summary of each antigen beside each entry.

□ CD molecule	Alternate Names	Locus ID	Past Guides
CD1a	R4; HTA1	909	CD1a
CD1b	R1	910	CD1b
CD1c	M241; R7	911	CD1c
CD1d	R3	912	CD1d
CD1e	R2	913	CD1e
CD2	CD2R; E-rosette receptor; T11; LFA-2	914	CD2
CD3delta	CD3d	915	
CD3epsilon	CD3e	916	
CD3gamma	CD3g	917	
CD4	L3T4; W3/25	920	CD4
CD5	Leu-1; Ly-1; T1; Tp67	921	CD5
CD6	T12	923	CD6
CD7	gp40	924	
CD8alpha	Leu2; Lyt2; T cell co-receptor; T8	925	
CD8beta	Leu2; CD8; Lyt3	926	
CD9	DRAP-27; MRP-1; p24	928	CD9
CD10	EC 3.4.24.11; neprilysin; CALLA; enkephalinase; gp100; NEP	4311	
CD11a	AlphaL integrin chain; LFA-1alpha	3683	CD11a
CD11b	AlphaM integrin chain; AlphaM-beta2; C3biR; CR3; Mac-1; Mo1	3684	CD11b
CD11c	AlphaX integrin chain; Axb2; CR4; leukocyte surface antigen p150,95	3687	CD11c
CDw12	p90-120	23444	CDw12
CD13	APN; EC 3.4.11.2; gp150	290	CD13
CD14	LPS-R	929	CD14
CD15u	Sulphated CD15		
CD16a	FCRIIIA	2214	
CD16b	FCRIIIB	2215	
CDw17	LacCer		CDw17

<input type="checkbox"/> CD molecule	Alternate Names	Locus ID	Past Guides
CD18	CD11a beta subunit; CD11b beta subunit; CD11c beta subunit; beta-2 integrin chain	<u>3689</u>	<u>CD18</u>
CD19	B4	<u>930</u>	<u>CD19</u>
CD20	B1; Bp35	<u>931</u>	
CD21	C3d receptor; CR2; EBV-R	<u>1380</u>	<u>CD21</u>
CD22	BL-CAM; Lyb8	<u>933</u>	<u>CD22</u>
CD23	B6; BLAST-2; FcεRII; Leu-20; Low affinity IgE receptor	<u>2208</u>	<u>CD23</u>
CD24	BA-1; HSA	<u>934</u>	<u>CD24</u>
CD25	IL-2R alpha chain; IL-2R; Tac antigen	<u>3559</u>	<u>CD25</u>
CD26	EC 3.4.14.5; ADA-binding protein; DPP IV ectoenzyme	<u>1803</u>	<u>CD26</u>
CD27	S152; T14	<u>939</u>	<u>CD27</u>
CD28	T44; Tp44	<u>940</u>	<u>CD28</u>
CD29	Platelet GPIIa; VLA-beta chain; beta-1 integrin chain	<u>3688</u>	
CD30	Ber-H2 antigen; Ki-1 antigen	<u>943</u>	<u>CD30</u>
CD31	GPIIa'; endocam; PECAM-1	<u>5175</u>	<u>CD31</u>
CD32	FCR II; Fc gamma RII	<u>2212</u>	
CD33	gp67; p67	<u>945</u>	<u>CD33</u>
CD34	gp105-120	<u>947</u>	<u>CD34</u>
CD35	C3bR; C4bR; CR1; Immune Adherence Receptor	<u>1378</u>	<u>CD35</u>
CD36	GPIIIb; GPIV; OKM5-antigen; PASIV	<u>948</u>	<u>CD36</u>
CD37	gp52-40	<u>951</u>	<u>CD37</u>
CD38	T10; cyclic ADP-ribose hydrolase	<u>952</u>	<u>CD38</u>
CD39		<u>953</u>	
CD40	Bp50	<u>958</u>	<u>CD40</u>
CD41	GPIIb; alpha IIb integrin chain	<u>3674</u>	
CD42a	GPIX	<u>2815</u>	<u>CD42a</u>

<input type="checkbox"/> CD molecule	Alternate Names	Locus ID	Past Guides
CD42b	GPIb α ; Glycocalicin	<u>2811</u>	<u>CD42b</u>
CD42c	GPIb-beta	<u>2812</u>	<u>CD42c</u>
CD42d	GPV	<u>2814</u>	<u>CD42d</u>
CD43	gpL115; leukocyte sialoglycoprotein; leukosialin; sialophorin	<u>6693</u>	<u>CD43</u>
CD44	ECMR III; H-CAM; HUTCH-1; Hermes; Lu, In-related; Pgp-1; gp85	<u>960</u>	<u>CD44</u>
CD44R	CD44v; CD44v9	<u>960</u>	<u>CD44R</u>
CD45	B220; CD45R; CD45RA; CD45RB; CD45RC; CD45RO; EC 3.1.3.4; LCA; T200; Ly5	<u>5788</u>	<u>CD45</u>
CD46	MCP	<u>4179</u>	<u>CD46</u>
CD47R	Rh-associated protein; gp42; IAP; neurophilin; OA3; MEM-133; formerly CDw149	<u>961</u>	<u>CD47</u>
CD48	BCM1; Blast-1; Hu Lym3; OX-45	<u>962</u>	<u>CD48</u>
CD49a	Alpha-1 integrin chain; VLA-1 alpha chain	<u>3672</u>	
CD49b	Alpha-2 integrin chain; GPIa; VLA-2 alpha chain	<u>3673</u>	
CD49c	Alpha-3 integrin chain; VLA-3 alpha chain	<u>3675</u>	
CD49d	Alpha-4 integrin chain; VLA-4 alpha chain	<u>3676</u>	<u>CD49d</u>
CD49e	Alpha-5 integrin chain; FNR alpha chain; VLA-5 alpha chain	<u>3678</u>	
CD49f	Alpha-6 integrin chain; Platelet gpI; VLA-6 alpha chain	<u>3655</u>	
CD50	ICAM-3	<u>3385</u>	<u>CD50</u>
CD51	VNR-alpha chain; alpha V integrin chain; vitronectin receptor	<u>3685</u>	
CD52		<u>1043</u>	<u>CD52</u>
CD53		<u>963</u>	<u>CD53</u>

<input type="checkbox"/> CD molecule	Alternate Names	Locus ID	Past Guides
CD54	ICAM-1	<u>3383</u>	<u>CD54</u>
CD55	DAF	<u>1604</u>	<u>CD55</u>
CD56	Leu-19; NKH1; NCAM	<u>4684</u>	<u>CD56</u>
CD57	HNK1; Leu-7	<u>964</u>	
CD58	LFA-3	<u>965</u>	<u>CD58</u>
CD59	1F-5Ag; H19; HRF20; MAC1F; M1RL; P-18; Protectin	<u>966</u>	<u>CD59</u>
CD60a	GD3		<u>CDw60</u>
CD60b	9-O-acetyl-GD3		<u>CDw60</u>
CD60c	7-O-acetyl-GD3		<u>CDw60</u>
CD61	CD61A; GPIIb/IIIa; beta 3 integrin chain	<u>3690</u>	
CD62E	E-selectin; ELAM-1; LECAM-2	<u>6401</u>	<u>CD62E</u>
CD62L	L-selectin; LAM-1; LECAM-1; Leu-8; MEL-14; TQ-1	<u>6402</u>	<u>CD62L</u>
CD62P	P-selectin; GMP-140; PADGEM	<u>6403</u>	<u>CD62P</u>
CD63	LIMP; MLA1; PTLGP40; gp55; granulophysin; LAMP-3; ME491; NGA	<u>967</u>	<u>CD63</u>
CD64	FC gammaRI; FCR I	<u>2209</u>	<u>CD64</u>
CD65	Ceramide-dodecasaccharide; VIM-2		
CD65s	Sialylated-CD65; VIM2		
CD66a	NCA-160; BGP	<u>634</u>	<u>CD66a</u>
CD66b	CD67; CGM6; NCA-95	<u>1088</u>	<u>CD66b</u>
CD66c	NCA; NCA-50/90	<u>4680</u>	<u>CD66c</u>
CD66d	CGM1	<u>1084</u>	<u>CD66d</u>
CD66e	CEA	<u>1048</u>	<u>CD66e</u>
CD66f	Pregnancy specific b1 glycoprotein; SP-1; PSG	<u>5669</u>	<u>CD66f</u>
CD68	gp110; macrosialin	<u>968</u>	<u>CD68</u>
CD69	AIM; EA 1; MLR3; gp34/28; VEA	<u>969</u>	<u>CD69</u>
CD70	CD27-ligand; Ki-24 antigen	<u>970</u>	

<input type="checkbox"/> CD molecule	Alternate Names	Locus ID	Past Guides
CD71	T9; transferrin receptor	<u>7037</u>	<u>CD71</u>
CD72	Ly-19.2; Ly-32.2; Lyb-2	<u>971</u>	
CD73	Ecto-5'-nucleotidase	<u>4907</u>	<u>CD73</u>
CD74	Class II-specific chaperone; Ii; Invariant chain	<u>972</u>	<u>CD74</u>
CD75	Lactosamines		
CD75s	Alpha-2,6-sialylated lactosamines (formerly CDw75 and CDw76)		<u>CDw75</u> ; <u>CDw76</u>
CD77	Pk blood group antigen; BLA; CTH; Gb3		<u>CD77</u>
CD79a	Ig alpha; MB1	<u>973</u>	
CD79b	B29; Ig beta	<u>974</u>	
CD80	B7; BB1	<u>941</u>	<u>CD80</u>
CD81	TAPA-1	<u>975</u>	<u>CD81</u>
CD82	4F9; C33; IA4; KAI1; R2	<u>3732</u>	<u>CD82</u>
CD83	HB15	<u>9308</u>	<u>CD83</u>
CD84		<u>8832</u>	<u>CD84</u>
CD85	ILT/LIR family {Young NT/Parham P.2001.IMMUN} {Allan DS/Braud VM.2000.IMMUN}	<u>10859</u>	<u>CD85</u>
CD86	B7-2; B70	<u>942</u>	<u>CD86</u>
CD87	uPAR	<u>5329</u>	<u>CD87</u>
CD88	C5aR	<u>728</u>	<u>CD88</u>
CD89	Fcalpha-R; IgA Fc receptor; IgA receptor	<u>2204</u>	<u>CD89</u>
CD90	Thy-1	<u>7070</u>	<u>CD90</u>
CD91	ALPHA2M-R; LRP	<u>4035</u>	
CD92	CTL1; formerly CDw92	<u>23446</u>	<u>CD92</u>
CDw93		<u>23447</u>	<u>CDw93</u>
CD94	Kp43	<u>3824</u>	<u>CD94</u>
CD95	APO-1; Fas; TNFRSF6; APT1	<u>355</u>	<u>CD95</u>
CD96	TACTILE	<u>10225</u>	

<input type="checkbox"/> CD molecule	Alternate Names	Locus ID	Past Guides
CD97		976	CD97
CD98	4F2; FRP-1; RL-388	4198	CD98
CD99	CD99R; E2; MIC2 gene product	4267	
CD100	SEMA4D	10507	CD100
CD101	IGSF2; P126; V7	9398	CD101
CD102	ICAM-2	3384	CD102
CD103	ITGAE; HML-1; integrin alphaE chain	3682	CD103
CD104	beta 4 integrin chain; TSP-1180; beta 4	3691	
CD105	endoglin	2022	CD105
CD106	INCAM-110; VCAM-1	7412	
CD107a	LAMP-1	3916	CD107a
CD107b	LAMP-2	3920	CD107b
CD108	SEMA7A; JMH human blood group antigen; formerly CDw108	8482	CD108
CD109	8A3; E123; 7D1		
CD110	MPL; TPO-R; C-MPL	4352	
CD111	PVRL1; PRR1; HevC; nectin-1; HIgR	5818	
CD112	HVEB; PRR2; PVRL2; nectin 2	5819	
CD113	Reserved		
CD114	CSF3R; HG-CSFR; G-CSFR	1441	CD114
CD115	c-fms; CSF-1R; M-CSFR	1436	
CD116	GM-CSF receptor alpha chain	1438	CD116
CD117	c-KIT; SCFR	3815	CD117
CD118	Reserved		
CDw119	IFNgR; IFNgRa	3459	
CD120a	TNFR1; p55	7132	
CD120b	TNFR2; p75; TNFR p80	7133	
CD121a	IL-1R; type 1 IL-1R	3554	
CDw121b	IL-1R, type 2	7850	

<input type="checkbox"/> CD molecule	Alternate Names	Locus ID	Past Guides
CD122	IL-2Rbeta	<u>3560</u>	<u>CD122</u>
CD123	IL-3Ralpha	<u>3563</u>	
CD124	IL-4R	<u>3566</u>	<u>CD124</u>
CDw125	IL-5Ralpha	<u>3568</u>	<u>CDw125</u>
CD126	IL-6R	<u>3570</u>	<u>CD126</u>
CD127	IL-7R; IL-7R alpha; p90 Il7 R	<u>3575</u>	<u>CD127</u>
CDw128a	CXCR1; IL-8RA	<u>3577</u>	
CDw128b	CXCR2; IL-8RB	<u>3579</u>	
CD129	Reserved		
CD130	gp130	<u>3572</u>	<u>CD130</u>
CD131	common beta subunit	<u>1439</u>	<u>CDw131</u>
CD132	IL2RG; common cytokine receptor gamma chain; common gamma chain	<u>3561</u>	<u>CD132</u>
<u>CD133</u>	PROML1; AC133; hematopoietic stem cell antigen; prominin-like 1	<u>8842</u>	
CD134	OX40	<u>7293</u>	
CD135	flt3; Flk-2; STK-1	<u>2322</u>	<u>CD135</u>
CDw136	msp receptor; ron; p158-ron	<u>4486</u>	<u>CDw136</u>
CDw137	4-1BB; ILA	<u>3604</u>	<u>CDw137</u>
CD138	heparan sulfate proteoglycan; syndecan-1	<u>6382</u>	
CD139		<u>23448</u>	<u>CD139</u>
CD140a	PDGF-R; PDGFRA	<u>5156</u>	
CD140b	PDGFRb	<u>5159</u>	
CD141	fetomodulin; TM	<u>7056</u>	<u>CD141</u>
CD142	F3; coagulation Factor III; thromboplastin; TF	<u>2152</u>	<u>CD142</u>
CD143	EC 3.4.15.1; ACE; kininase II; peptidyl dipeptidase A	<u>1636</u>	<u>CD143</u>
CD144	cadherin-5; VE-Cadherin	<u>1003</u>	<u>CD144</u>
CDw145			

<input type="checkbox"/> CD molecule	Alternate Names	Locus ID	Past Guides
CD146	MCAM; A32; MUC18; Mel-CAM; S-endo	<u>4162</u>	<u>CD146</u>
CD147	SA11; Basigin; CE9; HT7; M6; Neurothelin; OX-47; EMMPRIN; gp42	<u>682</u>	<u>CD147</u>
CD148	HPTP-eta; DEP-1; p260	<u>5795</u>	<u>CD148</u>
CDw149	new designation is <u>CD47R</u>		
CD150	SLAM; IPO-3; formerly CDw150	<u>6504</u>	<u>CDw150</u>
CD151	PETA-3; SFA-1	<u>977</u>	<u>CD151</u>
CD152	CTLA-4	<u>1493</u>	<u>CD152</u>
CD153	CD30L	<u>944</u>	
CD154	CD40L; T-BAM; TRAP; gp39	<u>959</u>	
CD155	PVR	<u>5817</u>	
CD156a	ADAM8; MS2 human; formerly CD156	<u>101</u>	<u>CD156a</u>
CD156b	ADAM17; TACE; cSVP	<u>6868</u>	
CD157	BP-3/IF-7; BST-1; Mo5	<u>683</u>	<u>CD157</u>
CD158	<u>KIR family</u> (detailed nomenclature to be published)		<u>KIR Family</u>
CD159a	NKG2A	<u>3821</u>	
CD160	BY55 antigen; NK1; NK28	<u>11126</u>	
CD161	KLRB1; NKR-P1A; killer cell lectin-like receptor subfamily B, member 1	<u>3820</u>	<u>CD161</u>
CD162	PSGL-1, PSGL	<u>6404</u>	<u>CD162</u>
CD162R	PEN5 (a post-translational modification of PSGL-1)	<u>6404</u>	
CD163	GHI/61; M130; RM3/1	<u>9332</u>	
CD164	MUC-24; MGC-24v	<u>8763</u>	
CD165	AD2; gp37	<u>23449</u>	<u>CD165</u>
CD166	BEN; DM-GRASP; KG-CAM; Neurolin; SC-1; ALCAM	<u>214</u>	<u>CD166</u>
<u>CD167a</u>	trkE; trk6; cak; eddr1; DDR1; MCK10; RTK6; NTRK4	<u>780</u>	
<u>CD168</u>	HMMR; IHABP; RHAMM	<u>3161</u>	

CD molecule	Alternate Names	Locus ID	Past Guides
CD169	sialoadhesin; siglec-1	6614	
CD170	Siglec-5	8778	
CD171	L1; L1CAM; N-CAM L1	3897	
CD172a	SIRP alpha	8194	
CD173	Blood group H type 2		
CD174	Lewis y	2525	
CD175	Tn		
CD175s	Sialyl-Tn		
CD176	TF		
CD177	NB1		
CD178	fas-L; TNFSF6; APT1LG1; CD95-L	356	
CD179a	VpreB; VPRED1; IGVPB	7441	
CD179b	IGLL1; lambda5; immunoglobulin omega polypeptide; IGVPB; 14.1 chain	3543	
CD180	LY64; RP105	4064	
CD183	CXCR3; GPR9; CKR-L2; IP10-R; Mig-R	2833	
CD184	CXCR4; fusin; LESTR; NPY3R; HM89; FB22	7852	
CD195	CCR5	1234	
CDw197	CCR7	1236	
CD200	OX2	4345	
CD201	EPC R	10544	
CD202b	tie2; tek	7010	
CD203c	NPP3; PDNP3; PD-Ibeta; B10; gp130RB13-6; ENPP3; bovine intestinal phosphodiesterase	5169	
CD204	macrophage scavenger R	4481	
CD205	DEC205	4065	
CD206	MRC1; MMR	4360	
CD207	Langerin	50489	

CD molecule	Alternate Names	Locus ID	Past Guides
CD208	DC-LAMP	<u>27074</u>	
CD209	DC-SIGN	<u>30385</u>	
CDw210	IL-10 R	<u>3587</u> ; <u>3588</u>	
CD212	IL-12 R	<u>3594</u>	
CD213a1	IL-13 R alpha 1	<u>3597</u>	
CD213a2	IL-13 R alpha 2	<u>3598</u>	
CDw217	IL-17 R	<u>23765</u>	
CD220	Insulin R	<u>3643</u>	
CD221	IGF1 R	<u>3480</u>	
CD222	Mannose-6-phosphate/IGF2 R	<u>3482</u>	
CD223	LAG-3	<u>3902</u>	
CD224	GGT; EC2.3.2.2	<u>2678</u>	
CD225	Leu13	<u>8519</u>	
CD226	DNAM-1; PTA1; TLI SA1	<u>10666</u>	
CD227	MUC1; episialin; PUM; PEM; EMA; DF3 antigen; H23 antigen	<u>4582</u>	
CD228	melanotransferrin	<u>4241</u>	
CD229	Ly9	<u>4063</u>	
CD230	Prion protein	<u>5621</u>	
CD231	TM4SF2; A15; TALLA-1; MXS1; CCG-B7; TALLA	<u>7102</u>	
CD232	VESP R	<u>10154</u>	
CD233	band 3; erythrocyte membrane protein band 3; AE1; SLC4A1; Diego blood group; EPB3	<u>6521</u>	
CD234	Fy-glycoprotein; Duffy antigen	<u>2532</u>	
CD235a	Glycophorin A	<u>2993</u>	
CD235b	Glycophorin B	<u>2994</u>	
CD235ab	Glycophorin A/B crossreactive mabs		
CD236	Glycophorin C/D		

29

<input type="checkbox"/> CD molecule	Alternate Names	Locus ID	Past Guides
CD236R	Glycophorin C	<u>2995</u>	
CD238	Kell	<u>3792</u>	
CD239	B-CAM	<u>4059</u>	
CD240CE	Rh30CE	<u>6006</u>	
CD240D	Rh30D	<u>6007</u>	
CD240DCE	Rh30D/CE crossreactive mabs		
CD241	RhAg	<u>6005</u>	
CD242	ICAM-4	<u>3386</u>	
CD243	MDR-1	<u>5243</u>	
CD244	2B4; NAIL; p38	<u>51744</u>	
CD245	p220/240		
CD246	Anaplastic lymphoma kinase	<u>238</u>	
CD247	Zeta chain	<u>919</u>	

Revised 07/02/02

prow@ncbi.nlm.nih.gov

The expression "the group of CD1 to CD247" as used herein means all the CD molecules in the list above.

5 In the most preferred embodiment, the anti-lymphoma antibody is directed against CD19, CD20, CD22, CD 30, in particular CD 20.

In the present patent application, an immunotargeting agent (immunoconjugate) is an agent carrying a cytotoxic moiety that, contrary to common cytotoxic medical agents, binds specifically to lymphatic tumor cell with a high affinity and which could be administered parentally, preferably intravenously, to a human being. In a preferred application, the immunotargeting agents are antibodies, which could be of different isotypes and could originate from any species. Of particular interest are the monoclonal antibodies and derivatives thereof. The latter include fragments such as the F(ab')₂, F(ab'), F(ab) and the like. They also include genetically engineered

hybrids or chemically synthesized peptides based on the specificity of the antigen binding region of one or several target specific monoclonal antibodies, e.g. chimeric or humanized antibodies, single chain antibodies etc.

The biomolecule binding moiety, which is an anti-lymphoma antibody reactive moiety, is bound or conjugated to the anti-lymphoma antibody, either covalently or non-covalently with an affinity binding constant of at least 10^8M^{-1} .

The term "anti-lymphoma antibody" used herein is intended to mean an antibody with the ability of specific binding to a CD antigen on lymphoma tumour cells with an affinity binding constant of at least $5 \times 10^6 \text{M}^{-1}$, preferably at least 10^8M^{-1} .

The term "variants" of the anti-lymphoma antibody as used herein means any modifications, fragments or derivatives thereof having the same or essentially similar affinity binding constant when binding to the CD antigen molecule, i.e. an affinity binding constant of at least $5 \times 10^6 \text{M}^{-1}$, preferably at least 10^8M^{-1} .

Any of these variants could have been modified by the coupling of various number of polyethylene glycol chains in order to optimise the half-life in body fluid and the retention of the antibody or antibody fragments or derivatives, in the tumor tissue. In the most preferred application, the antibodies or antibody derivatives should allow for the attachment of a sufficient number of biotin residues to be used for extracorporeal removal through interaction with immobilized avidin, without significantly diminishing the binding properties of the targeting agent.

In order to enhance the specificity, tumour specific monoclonal antibodies are used as a carrier (immunoconjugates) of various cytotoxic moieties, such as, but not limited to, radio nuclides, chemotherapy drugs, synthetic or natural occurring toxins, immunosuppressive agents,

immunostimulating agents and enzymes used in pro-drug protocols. The cytotoxic moiety is preferably a radionuclide such as a gamma-emitter e.g. iodine-131 or metal ion conjugate, where the metal is selected from a beta-particle emitter, such as yttrium or rhenium. US Patent No. 4,472,509, Gansow, et al., discloses the use of diethylenetriaminepentaacetic acid (DTPA) chelating agents for the binding of radio metals to monoclonal antibodies. The patent is particularly directed to a purification technique for the removal of non-bonded and adventitious-ly bonded (non-chelated) metal from radiopharmaceuticals but is illustrative of art recognized protocols for preparation of radionuclide labelled antibodies.

According to such general procedures, an antibody specifically reactive with the target tissue associated antigen is reacted with a certain quantity of a selected bifunctional chelating agent having protein binding and metal binding functionalities to produce a chelator/-antibody conjugate. In conjugating the antibodies with the chelators, an excess of chelating agent is reacted with the antibodies, the specific ratio being dependent upon the nature of the reagents and the desired number of chelating agents per antibody. It is a requirement that the radionuclides be bound by chelation (for metals) or covalent bonds in such a manner that they do not become separated from the biotinylation/radiolabeling compound under the conditions that the biomolecule conjugates is used (e.g. in patients). Thus, the most stable chelates or covalent bonding arrangements are preferred. Examples of such binding/bonding moieties are: aryl halides and vinyl halides for radionuclides of halogens; N_2S_2 and N_3S chelates for Tc and Re radionuclides; amino-carboxy derivatives such as EDTA, DTPA, derivatives of Me-DTPA and Cyclohexyl-DTPA, and cyclic amines such as NOTA, DOTA, TETA, CITC-DTPA, and triethylenetetraaminehexa-acetic acid derivatives (Yuangfang and Chuanchu, Pure & Appl. Chem. 63, 427-463, 1991) for In, Y, Pb, Bi, Cu, Sm,

and Lu radionuclides, and where the radionuclide is, but not limited to, any of the following elements:

Beta radiation emitters, which are useful as cytotoxic agents, include isotopes such as scandium-46, scandium-47, scandium-48, copper-67, gallium-72, gallium-73, yttrium-90, ruthenium-97, palladium-100, rhodium-101, palladium-109, samarium-153, lutetium-177, rhenium-186, rhenium-188, rhenium-189, gold-198, radium-212 and lead-212. The most useful gamma emitters are iodine-131 and indium-114. Other metal ions useful with the invention include alpha radiation emitting materials such as 212-bismuth, 213-bismuth, and At-211 as well as positron emitters such as gallium-68 and zirconium-89.

In another embodiment of the invention, radionuclide-labelled targeting agents are useful not only in the treatment of lymphatic cancers, but also for imaging of such cancers.

At a suitable time after administration, "cytotoxic targeting agents" will be cleared from the blood system by extracorporeal means. To facilitate the extracorporeal depletion, an apparatus for extracorporeal circulation of whole blood or plasma will be connected to the patient through tubing lines and blood access device(s). Such an apparatus should provide conduits for transporting the blood to an adsorption device and conduits for returning the processed blood or plasma to the patient. In the case plasma is processed through the adsorption device, a plasma separation device is needed as well as means of mixing the concentrated blood with processed plasma. The latter is normally achieved by leading the two components into an air-trap where the mixing occurs.

In the case where whole blood is processed, an ordinary dialysis machine can constitute the base for such an apparatus. Dialysis machines are normally equipped with all the necessary safeguards and monitoring devices to meet patient safety requirements as well as to allow easy handling of the system. Hence, in a preferred embodiment

whole blood is processed and a standard dialysis machine is utilised with only minor modifications of the hardware. However, such a machine requires a new program fitted to the newly intended purpose.

5 In addition to the apparatus, special blood line tubings suitable for the intended flow and distance from the patient and the machine are needed. These line tubings could be made of any material compatible with blood or plasma and would include materials used in ordinary
10 tubings used in dialysis.

Blood access could be achieved through peripheral vein catheters or, if higher blood flow is needed, through central vein catheters such as, but not limited to, subclavian or femoral catheters.


15 For affinity adsorbents, the matrix may be of various shape and chemical composition. It may, for example, constitute a column house filled with particulate polymers, the latter of natural origin or artificially made. The particles may be macroporous or their surface may be
20 grafted, the latter in order to enlarge the surface area. The particles may be spherical or granulated and be based on polysaccharides, ceramic material, glass, silica, plastic, or any combination of these or alike materials. A combination of these could, for example, be solid par-
25 ticles coated with a suitable polymer of natural origin or artificially made. Artificial membranes may also be used. These may be flat sheet membranes made of cellulose, polyamide, polysulfone, polypropylene or other types of material which are sufficiently inert, biocompatible, non-
30 toxic and to which the receptor could be immobilized either directly or after chemical modification of the membrane surface. Capillary membranes like the hollow fibers made from cellulose, polypropylene or other materials suitable for this type of membranes may also be used. A
35 preferred embodiment is a particulate material based on agarose and suitable for extracorporeal applications.

In one embodiment, an affinity label is attached to the anti-lymphoma antibody and the adsorption device contains an immobilized receptor binding specifically to the affinity ligand. Any type of affinity ligand/immobilized receptor combinations such as "antibodies and antigens/haptens" and "protein and co-factors" could be used in the this application, provided that they exhibit a sufficiently high binding affinity and selectively to the tumor markers, and that the affinity ligand-receptor interaction is not interfered with by blood or other body fluids or tissues being in contact with the immunotargeting agent and/or the device.


In one of the most preferred applications, the affinity ligand/immobilized receptor combination is biotin or biotin derivatives and biotin binding molecules, in particular where the affinity ligand is biotin or derivatives thereof and the immobilized receptor is avidin or streptavidin or any other biotin binding molecule. The affinity ligand pairs of biotin/avidin and biotin/streptavidin are often used with biomolecules. The very strong interaction (i.e. $K = 10^{13} - 10^{15} \text{ M}^{-1}$) of biotin with the proteins avidin and streptavidin (Green, Methods Enzymol. 184, 51-67, 1990; Green, Adv. Prot. Chem. 29, 85-133, 1975) provides a foundation for their use in a large number of applications, both for *in vitro* and *in vivo* uses. A further application of the invention is the simultaneous removal of several different biotinylated "anti-cancer agents" through the same extracorporeal procedure.

The reagent used in the present invention is schematically shown below, wherein the biomolecule reactive moiety is an anti-lymphoma reactive moiety.

35



10 The medical agent according to the present invention is schematically shown below, wherein an anti-lymphoma antibody is bound or conjugated to the reagent via the anti-lymphoma antibody reactive moiety of the reagent.



25 In the schematically shown reagent and medical agent, respectively, the different components will be presented in more detail below.

The anti-lymphoma antibody reactive moiety is chosen from a group of active esters consisting of N-hydroxy-succinimide esters, sulfo-N-hydroxysuccinimide esters, and phenolic esters; aryl and alkyl imidates; alkyl or aryl isocyanates or isothiocyanates reacting with amino groups on the anti-lymphoma antibody, or maleimides or alpha-haloamides reacting with sulfhydryl groups on the anti-lymphoma antibody; or aryl or alkylhydrazines or alkyl or arylhydroxylamines reacting with aldehyde or

30

35

ketone groups naturally occurring or synthetically produced on the anti-lymphoma antibody, or variants thereof.

The effector agent is a radionuclide binding moiety, optionally provided with a radionuclide, a synthetic or naturally occurring toxin, an enzyme capable of converting pro-drugs to active drugs, immunosuppressive or immunostimulating agents, radiosensitizers, enhancers for X-ray or MRI or ultrasound, non-radioactive elements, which can be converted to radio active elements by means of external irradiation after that the anti-lymphoma antibody carrying said element has been accumulated to specific cells or tissues, or photoactive compounds or compounds used in photo imaging or photo dynamic therapy, or any other molecule having the same or similar effect, directly or indirectly, on lymphoma cells or lymphoma tissues. More precisely, the effector agent comprises Me-DTPA, CITC-DTPA, and cyclohexyl-DTPA.

The affinity ligand can be any moiety that binds with another molecule with an affinity constant of 10^6 M^{-1} or higher. A preferred affinity ligand is a moiety which binds specifically to avidin, streptavidin, or any other derivatives, mutants or fragments of avidin or streptavidin having essentially the same binding function to the affinity ligand. Preferably, the affinity ligand is biotin, or a biotin derivative having essentially the same binding function to avidin or streptavidin as biotin. Said biotin derivative may be chosen from the group consisting of a biotin derivative having essentially the same binding function to avidin or streptavidin as biotin.

The anti-lymphoma antibody having ability to be conjugated to said anti-lymphoma antibody reactive moiety interacts with one or more different cell surface antigen(s) present on the surface of lymphoma tumour cells, said one or more cell surface antigen(s) being one or more different CD antigen(s), or variants thereof, wherein the anti-lymphoma antibody preferably is chosen

from anti-CD20 antibodies, preferably rituximab, ibritumomab and tositumomab.


The trifunctional cross-linking moiety is chosen from the group consisting of triaminobenzene, tricarboxy-
5 benzene, dicarboxyanyline and diaminobenzoic acid.

Linker 1 is a chemical moiety that is an attaching moiety and spacer between the trifunctional cross-linking moiety and the affinity ligand, preferably a biotin moiety, such that binding with avidin or streptavidin, or
10 any other biotin binding species, is not diminished by steric hindrance. Linker 1 may also impart increased water solubility and biotinidase stabilization, preferably against cleavage by biotinidase by introduction of an alpha carboxylate or an N-methyl group. Further, it
15 contains hydrogen bonding atoms, preferably ethers or tioethers, or ionisable groups, preferably carboxylate, sulfonates, or ammonium groups to aid in water solubilisation of the biotin moiety.

For the structural requirements of the biotin containing moiety, the following applies with reference to
20 the following embodiment of the present invention:

generalized structure of a 1033-anti-CD20 antibody

25



This structure is bound to the effector agent, wherein the anti-CD20 antibody preferably is rituximab, wherein n is 2-4, preferably 3; o is 1-6, preferably 3; p is 1-6, preferably 3; R_2 is $=CH_2OH$ or $-CO_2H$; and R_1 is -
5 CH_3 , $-CH_2OH$, or $-H$.

There are three aspects of the biotin portion of the 1033 structures that are important in this application: (1) blockage of biotinidase cleavage, (2) retention of high biotin binding affinity, and (3) attainment of a
10 reasonable aqueous solubility. To provide these attributes, biotin conjugates must be composed of a biotin molecule and an appropriate linker, which are coupled to a cross-linking moiety.

Biotin conjugates must be prepared by conjugation
15 with the carboxylate on the pentanoic acid side chain ($n = 3$). Conjugation at other locations in the biotin molecule results in complete loss of binding with avidin and streptavidin. This renders the biotin molecule useless for this application. The preferred form of conjugation
20 is formation of an amide bond with the carboxylate group (as depicted in the general formula). Since binding of biotin with avidin and streptavidin is in a deep pocket (e.g. 9Å), shortening ($n < 3$) or lengthening ($n > 3$) of the pentanoic acid side chain results in low binding affinity,
25 ty, which is not desired for this application.

Blocking of biotinidase activity is achieved by attaching appropriate substituents to the biotinamide amine (i.e. R_1) or to an atom adjacent to that amine (i.e. R_2). Biotinidase is an enzyme that cleaves (hydro-
30 lyzes) the amide bond of biotin carboxylate conjugates. This enzyme is very important in recycling biotin in animals and man. Metabolism of biotin in (several different) protein carboxylases releases biotin-w-N-lysine (biocytin), and biotinidase specifically cleaves that
35 amide bond to release free biotin. Biotinidase is also capable of cleaving (non-specifically) other biotinamide bonds. In this application, it is important that biotini-

dase do not cleave biotin from the conjugates, otherwise the desired outcome will not be achieved. Thus, the useful biotin conjugate structures incorporate functional groups (R_1 or R_2) that block the enzymatic activity of biotinidase. While it is likely that any structure for R_1 will block biotinidase, its structure is generally limited to a methyl (CH_3) group, as this group completely blocks biotinidase activity. The N-methyl group decreases the binding affinity of biotin with avidin and strept-
avidin significantly, but it is still useful in this application. Larger groups for R_1 (e.g. ethyl, aryl, etc.) are not useful due to the loss of binding affinity. The alternative to having a substituent R_1 is to have a substituent R_2 on the atom (e.g. methylene) adjacent to the biotinamide amine. Much larger and more varied substituents can be used in this position without any significant effect on the binding affinity of biotin. Biotinidase is not completely blocked when $R_2 = \text{CH}_3$ or CH_2CH_3 , although the rate of cleavage is slowed considerably (i.e. to 25% and 10% respectively). Complete blockage of biotinidase activity is attained when $R_2 = \text{CH}_2\text{OH}$ and CO_2H functionalities. The important consideration is that there is no decrease in binding affinity when these groups are incorporated as R_2 . Larger functional groups can also be used as R_2 to block biotinidase activity, but results in a decrease in binding affinity. The larger functional groups as R_2 are useful in this application if they do not cause a decrease in binding affinity greater than that obtained when $R_1 = \text{CH}_3$.

The biotin affinity and water solubility of the biotin moiety in 1033 are affected by the linker moiety used. The length and nature of the linker moiety (Linker 1) will be dependent to some degree on the nature of the molecule that it is conjugated with. The linker moiety serves the function of providing a spacer between the biotin moiety and the rest of the conjugate such that the biotin binding is not affected by steric hindrance from

the protein (or other conjugated molecule). The length (number of atoms in a linear chain) of the linker may vary from $n = 4-20$ for conjugates with small molecules (e.g. steroids) to $n > 20$ for large conjugate molecules (e.g. IgG molecule). The nature of the atoms in the linker (linear chain or branch from it) will also vary to increase water solubility. For example, linkers that contain more than 4 methylene units are improved by incorporation of oxygen or sulfur atoms (forming ethers or thioethers) or by having appended ionizable functionalities (e.g. sulfonates, carboxylates, amines or ammonium groups).


Linker 2, if present, is a chemical moiety that is used to attach the radionuclide binding moiety to the trifunctional cross-linking moiety. It provides a spacer length of 1-25 atoms, preferably a length of 6-18 atoms, or groups of atoms. Linker 2 may also impart increased water solubility due to the presence of hydrogen bonding atoms, preferably ethers or bioethers, or ionisable groups, to aid in water solubilisation.

Linker 3 may not be required, but where advantageous, it is a chemical moiety used to attach the biomolecule reactive moiety to the trifunctional cross-linking moiety. Linker 3 may be used as a spacer with a length of 1-25 atoms, preferably 6-18 atoms, or groups of atoms and/or it may be used to increase the water solubility of the compound due to the presence of hydrogen bonding atoms, such as ethers or thioethers, or ionisable groups, preferably carboxylate, sulfonates, or ammonium groups to aid in water solubilisation.

Moreover, the reagent according to the present invention may contain more than one affinity ligand and/or more than one effector agent bound to a trifunctional or tetrafunctional cross-linking moiety.

A preferred embodiment of the medical agent according to the present invention has the following schematic structure:


41



where the chelating group is, but not limited to, any of the following compounds: aryl halides and vinyl halides for radionuclides of halogens; N_2S_2 and N_3S chelates for Tc and Re radionuclides; amino-carboxy derivatives such as EDTA, DTPA, derivatives Me-DTPA and Cyclohexyl-DTPA, and cyclic amines such as NOTA, DOTA, TETA, CITC-DTPA, and triethylenetetraaminehexaacetic acid derivatives (Yuangfang and Chuanchu, Pure & Appl. Chem. 63, 427-463, 1991) for In, Y, Pb, Bi, Cu, Sm, Lu radionuclides and where the radionuclide is, but not limited, any of the following elements: Beta radiation emitters, which are useful as cytotoxic agents, include isotopes such as scandium-46, scandium-47, scandium-48, copper-67, gallium-72, gallium-73, yttrium-90, ruthenium-97, palladium-100, rhodium-101, palladium-109, samarium-153, lutetium-177, rhenium-186, rhenium-188, rhenium-189, gold-198, radium-212 and 212 lead. The most useful gamma emitters are iodine-131 and indium-114. Other metal ions useful with the invention include alpha radiation emitting materials such as 212-bismuth, 213-bismuth, and At-211 as well as positron emitters such as gallium-68 and zirconium-89.

In the most preferred embodiment of the present invention, the medical agent is the rituximab conjugate with 1-5 groups of 3-(13'-Thiourea benzylDOTA)Trioxadamine-1-(13''-Biotin-Asp-OH)Trioxadamine-5-isothiocy-

nato-Aminoisophthalate (see below). The radionuclide is ^{90}Y for therapeutic application and ^{111}In for in vivo diagnostic application. In the very most preferred embodiment, the rituximab conjugate contains 1.5 - 3.5 groups of 3-(13'-ThioureabenzylDOTA)Trioxdiamine-1-(13''-Biotin-Asp-OH)Trioxdiamine-5-isothiocyanato-Aminoisophthalate.



Ibritumomab or tositumomab is also effective as anti-lymphoma antibody in the medical agent.

Examples

The following examples shall not be construed as limiting the invention, but should be regarded as evidence of the applicability of the invention.

Example 1 - Conjugation and radiolabelling of rituximab.

In this and subsequent examples, Indium-111 has in some instances been used as a substitute for Yttrium-90, because the former is a gamma-emitter and possesses less radiation hazard than Yttrium-90.

The monoclonal antibody, Rituximab was conjugated with 3-(13'-ThioureabenzylDOTA)Trioxdiamine-1-(13''-Biotin-Asp-OH)trioxdiamine-5-Isothiocyanato-Aminoisophthalate (MitraTag-1033), for short also called "1033" in the following, using the method described by Wilbur D.S et al in Bioconjugate Chem. 13:1079-1092, 2002. A 5 mg quantity of the monoclonal antibody was dialysed against

1L metal free HEPES with a minimum of 5 buffer changes over 3 days at 4°C. A solution of MitraTag-1033 was made in water, and an appropriate volume was added to the antibody solution. After incubation overnight at room temperature, the antibody-conjugate was dialysed against 1L metal free 500 mM ammonium acetate buffer pH 5.3 with a minimum of 3 buffer changes over 3 days at 4°C. The demetalated conjugated antibody was stored at 4-8°C until used in radiolabelling experiments.

275 µl antibody conjugate (1375 µg; 1033-Rituximab) in 500 mM ammonium acetate buffer pH 5.3 was mixed with 15 µl ¹¹¹InCl₃ (or ⁹⁰YCl₃) in 50 mM HCl. The labelling was conducted at 45°C for 16 minutes. 28 µl DTPA was added to stop reaction. The quality of the radio conjugate was determined by TLC and HPLC. The number of MitraTag-1033 per monoclonal antibody molecule was determined by the HABA method.

Example 2 - Binding of the 1033-conjugated monoclonal antibody to an avidin-adsorbent.

The fraction of the 1033-rituximab radio conjugate binding to the Avidin-adsorbent utilised in the Mitradep[®] device, was analysed utilising micro-columns.

The non-bound protein fraction of a 2.4 conjugates/-IgG 1033-rituximab was 9%, and of a 4.6 conjugates/IgG 1033-rituximab 3%. This is well in line with a Poisson distribution of the conjugates. Hence, the above Rituximab conjugates should contain fractions, which are not labelled with MitraTag-1033. Hence, the non-binding fraction complies with the expected fraction of non-conjugated Rituximab i.e. the non-radioactive fraction.

More than 99% of the radioactivity in a radiolabelled 1033-conjugate sample was bound to the micro-column with the Avidin-adsorbent.

Example 3 - Depletion of 1033-rituximab conjugates during in vitro simulated treatments.

The depletion kinetics of 1033-rituximab during a patient treatment was simulated *in vitro* utilising a

recirculation method based on the principles described by Schindhelm K. (Artificial Organs 13:21-27 (1989)).

The 1033-rituximab was diluted in a solution with the same relative viscosity as human blood, and was re-circulated in vitro through a small-scale model of the Mitradep[®] device. 125 ml of a blood substitute, containing 10 mg of 1033-rituximab, were re-circulated at 6.25 ml/min (corresponds to 100 ml/min in Mitradep[®]). Three reservoir volumes were processed. The levels of 1033-rituximab in the reservoir were monitored.

When two preparations of 1033-rituximab with different numbers of MitraTag[™]-1033 moieties per Rituximab molecule were analysed, the results presented in Fig. 1 were obtained. As seen, the depletion of 1033-rituximab is not different from the theoretical depletion line, i.e. all 1033-rituximab present in the solution passing through the device is removed. Studies with biotinylated human IgG have shown that an efficient depletion is obtained at a biotin/IgG ratio down to 1.4 biotin/IgG (lowest ratio tested).

It was concluded that 1033-rituximab could be efficiently removed during an extracorporeal affinity adsorption procedure utilising the device Mitradep[®].

Example 4 - Binding of the 1033-conjugate to the target antigen CD20.

After conjugation with MitraTag[™]-1033, the 1033-rituximab conjugates were analysed for binding to the target antigen CD20 to confirm that the conjugation process has not denaturated the antigen binding. The CD20 antigen is not available in purified and soluble form. Therefore, during testing the CD20 expressing B-cell lymphoma cell lines, Raji and/or SB, was utilised as targets.

The specificity of the antigen binding was analysed by immunofluorescence in a flowcytometry (FACS) method. Briefly, the Raji cells were incubated with biotinylated-Rituximab and 1033-rituximab conjugates. After incuba-

tion, the cells were washed and incubated with fluorescence-labelled Avidin. After washing the cells were analysed in the FACS instrument. As positive control, a biotinylated mouse monoclonal antibody against CD20 was used, as negative control a PBS buffer was used. For control of binding to F_c-receptors on the cells, biotinylated normal human IgG was used. The results are presented in graphs where the x-axis presents the amount of fluorescence per cell on a logarithmic scale, and the Y-axis the number of cells displaying the specified fluorescence.

As seen in Fig. 2, no non-specific binding of Avidin to the cells was detected. Neither was binding to F_c-receptors seen utilising biotinylated human IgG. There was no significant difference in binding between the control mouse antibody, biotinylated Rituximab, or the two MitraTag™-1033 conjugates tested.

The specificity was also determined by analysing the binding of the conjugates to a CD20-positive cell line (SB) and a CD20-negative cell line (HSB) established from the same individual in an ELISA. The cells were dried into the wells of an ELISA plate. After incubation with 1033-rituximab conjugates, the bound antibodies were detected with an enzyme-conjugated Streptavidin. Biotinylated Rituximab and biotinylated normal human IgG were used as positive and negative control, respectively. As seen in Fig. 3, non-specific binding to the control cells was insignificant.

It was concluded that Rituximab retains the binding specificity to the antigen CD20 after conjugation with the MitraTag™-1033 reagent.

Example 5 - Analyses of the affinity of the binding to the CD20 antigen

The influence of the conjugation process on the binding affinity (strength) of Rituximab to the target antigen CD20 was studied utilising a competitive inhibition assay.

Briefly, increasing amounts of non-radiolabelled Rituximab and 1033-rituximab conjugates were mixed with a constant amount of ^{125}I -labelled Rituximab labelled utilising the Bolton-Hunter reagent. The mixtures were added to fixed SB lymphoma cells in 96 plate wells. After incubation for 2 hours at room temperature, the wells were washed, and the radioactivity bound to the cells was measured in an automatic NaI(Tl) scintillation well counter.

For each concentration of cold Rituximab and 1033-rituximab conjugates, the per cent inhibition of cell binding radioactivity was calculated. The per cent inhibition was plotted against concentration (Fig. 4), and the concentration required for 50% inhibition (IC_{50}) was calculated from the graph (Table 1). The IC_{50} is a measure of the relative affinity (avidity) of the tested antibody; a decrease of affinity is seen as an increased IC_{50} concentration. To be a significant change in affinity it is often stated that the difference in IC_{50} should be at least 10-fold.

Table 1		
Sample	IC_{50} (nM)	IC_{50} (relative rituximab)
Rituximab	26	1.0
1.6 1033-rituximab ⁽¹⁾	106	4.1
2.4 1033-rituximab	100	3.8
3.4 1033-rituximab	350	13.5
4.6 1033-rituximab	440	16.9
Human IgG	No inhibition	--

(1) 1.6 1033-rituximab denotes 1033-rituximab conjugated with 1.6 MitraTag/rituximab

A slight decrease in affinity was seen for the 1.6- and 2.4 1033-rituximab conjugates, whereas the decrease for the 3.4- and 4.6- 1033-rituximab conjugates was slightly above ten-fold relative to the IC50 concentration of Rituximab. The affinity for the 3.4- and 4.6- 1033-rituximab conjugates is probably still high enough to obtain a proper tumour uptake in patients.

It has been shown in clinical studies that a tenfold difference in affinity does not result in any significant difference in tumour uptake (ref. 6). Therefore, it was concluded that conjugation of Rituximab with up to 3-4 conjugates per antibody would not diminish the binding properties of the antibody *in vivo*.

Example 6 - Pharmacokinetics of MitraTag-1033 conjugates of rituximab.

Rats of the Sprague Dawley strain were injected intravenously with approximately 50 μ g of 1033-rituximab (4.6 1033 moieties per antibody) labelled with 3 - 4 MBq 111 Indium mixed with 1.2 mg/rat of a non-conjugated Rituximab. Whole body (WB) imaging was performed using a scintillation camera (General Electric 400T, GE, Milwaukee, WI, USA) equipped with a medium-energy collimator. Images were stored and analysed with Nuclear MAC 2.7 software. From images, the total number of counts in the entire body were obtained. After radioactivity decay correction and background subtraction, the counts were used for the calculation of activity retention (%) in the body. See Fig. 5.

To define pharmacokinetics of 111 In-1033-rituximab and compare it with 111 In-DOTA-hMn14, about 0.2 ml blood was obtained from the periorbital venous plexa on the following occasions: 10 min, 1, 8, 24, 48 and 96 hours post injection. The radioactivity was measured in an automatic NaI(Tl) scintillation well counter and expressed in percent of injected activity per gram blood (%/g) corrected for 111 In decay (Fig. 6).

Example 7 - Biodistribution of conjugates to organs and tissues.

At dissections, performed after 1, 8, 24, 48, and 96 hours post injection, organs and tissues of interest were removed, weighed and measured for radioactivity content. The radioactivity was measured in an automatic NaI(Tl) scintillation well counter, and the counts were corrected for decay. The distribution of the injected activity is shown in Fig. 7, and Table 2.

10

Table 2	Uptake of ¹¹¹ In-1033-rituximab (% injected dose/g)				
	1 h	8 h	24 h	48 h	96 h
Tissue					
Muscle	0.06	0.06	0.12	0.13	0.11
Kidney	0.98	0.91	0.83	0.86	1.00
Liver	1.22	1.79	1.86	1.92	3.42
Spleen	1.02	0.99	1.04	1.30	1.31
Bowel	0.10	0.29	0.32	0.29	0.20
Lymph nodes	0.26	1.03	1.99	2.88	2.54
Lung	0.74	0.89	0.71	0.52	0.38
Bone marrow	0.79	0.62	0.57	0.65	0.52

Example 8 - In vivo stability of the radiolabelled MitraTag-1033 antibody conjugates.

The stability of the MitraTag™-1033 moiety *in vivo* was determined by analysing the percentage of radioactivity in blood binding to Avidin-microcolumns.

About 0.1 ml blood was obtained from the periorbital venous plexa on following occasions: 1, 8, 24, 48, and 96 hours post injection. 50 µl blood was applied to a micro-column with Avidin-agarose (0.3 ml adsorbent). After incubation for 10 minutes, the unbound radioactivity was washed off the column. The radioactivity in the column and the collected washing fluid was measured in an automatic NaI(Tl) scintillation well counter and the bound fraction was expressed in percent of the total radioactivity applied to the column.

Time post injection (hours)	% Avidin-binding	Range (%)	Animals analysed
0	99.2	99.1 - 99.3	-
1	99.4	99.4 - 99.5	3
8	99.4	99.4 - 99.4	3
24	99.3	99.2 - 99.4	2
48	99.1	98.9 - 99.3	3
96	98.5	97.7 - 99.1	3

Sample 0 is on the conjugate to be injected.

During the period studied, no reduction of the binding to avidin of the radioactivity present in blood could be detected.

Therefore, it was concluded that the linkage between biotin and the DOTA chelate is stable in blood circulation up to 96 hours post injection.

When plasma was separated on a HPLC size exclusion column, no significant change in size distribution could be seen when a 10 min sample was compared with a 47 hour sample (Fig. 8).

Example 9 - Treatment regime in B-cell lymphoma according to the most preferred embodiment of the invention.

The treatment regime can be separated in the following events:

- All patients will receive a dose of 250 mg/m² rituximab one week prior to therapy (day -7) in order to eliminate the circulating B-cells, immediately

followed by a diagnostic dose of 50-150 MBq (1.5-4 mCi) ^{111}In -1033-rituximab.

- On day 0 all patients will receive 250 mg/m² rituximab immediately followed by a therapeutic dose of ^{90}Y -1033-rituximab (>10MBq/kg bodyweight). Patients may, optionally, be administered a dose of 150-250 MBq (4-7 mCi) ^{111}In -1033-rituximab, which will be used for imaging for dosimetry.
- On day 1 or 2, patients are treated with Mitradep[®], allowing 3 blood volumes to pass the Mitradep[®] device.

CLAIMS

1. A medical agent comprising a reagent conjugated to an anti-lymphoma antibody or a variant thereof, where-
5 in the reagent is a single molecule with at least three functional parts b) - d), wherein
- a) a trifunctional cross-linking moiety is coupled to
 - b) an affinity ligand via a linker 1, to
 - 10 c) an effector agent via a covalent bond, optionally via a linker 2, and to
 - d) a biomolecule reactive moiety, optionally via a linker 3, wherein said biomolecule reactive moiety is an anti-lymphoma antibody reactive moiety being
15 capable of forming a bond with the anti-lymphoma antibody or a variant thereof, thereby forming the conjugate, and wherein the anti-lymphoma antibody or variant thereof is interacting with one or more different CD antigens present on the surface of
20 lymphoma tumour cells.
2. The medical agent according to claim 1, wherein the anti-lymphoma antibody or variant thereof is mono-clonal and is interacting with one or more CD antigen(s) chosen from the group of CD1 to CD247.
- 25 3. The medical agent according to claim 2, wherein the anti-lymphoma antibody or variant thereof is interacting with CD19, CD20, CD22, and CD30, most preferably CD20.
- 30 4. The medical agent according to any one of the preceding claims, wherein the anti-lymphoma antibody is ibritumomab, rituximab, or tositumomab, preferably rituximab.
- 35 5. The medical agent according to any one of the preceding claims, wherein the anti-lymphoma antibody variant has the same or essentially the same ability as the anti-lymphoma antibody to bind to both the anti-lymphoma antibody reacting moiety and said CD antigen/-

antigens on the surface of lymphoma tumour cells, and wherein said variant is an antibody derivative, preferably the $F(ab')_2$, $F(ab')$, or $F(ab)$ fragment; genetically engineered hybrids or chemically synthesized peptides, preferably chimeric or humanized antibodies, and single chain antibodies.

6. The medical agent according to any one of the preceding claims, wherein the anti-lymphoma antibody or variant thereof binds to said cell surface antigen(s) on lymphoma tumour cells with an affinity binding constant of at least $5 \times 10^6 M^{-1}$, preferably at least $10^8 M^{-1}$.

7. The medical agent according to any one of the preceding claims, wherein the bond formed between the anti-lymphoma antibody reactive moiety and the anti-lymphoma antibody or a variant thereof is either covalent or non-covalent with a binding affinity constant of at least $10^8 M^{-1}$.

8. The medical agent according to claim 1, wherein the anti-lymphoma antibody reactive moiety is chosen from a group of active esters consisting of N-hydroxysuccinimide esters, sulfo-N-hydroxysuccinimide esters, and phenolic esters; aryl and alkyl imidates; alkyl or aryl isocyanates or isothiocyanates reacting with amino groups on the anti-lymphoma antibody, or maleimides or alpha-haloamides reacting with sulfhydryl groups on the anti-lymphoma antibody; or aryl or alkylhydrazines or alkyl or arylhydroxylamines reacting with aldehyde or ketone groups naturally occurring or synthetically produced on the anti-lymphoma antibody.

9. The medical agent according to claim 8, wherein said anti-lymphoma antibody reactive moiety also includes variants having essentially the same ability to bind to said anti-lymphoma antibody.

10. The medical agent according to claim 1, wherein the effector agent is a radionuclide binding moiety, optionally provided with a radionuclide, a synthetic or naturally occurring toxin, an enzyme capable of convert-

ing pro-drugs to active drugs, immunosuppressive or immunostimulating agents, radiosensitizers, enhancers for X-ray or MRI or ultrasound, non-radioactive elements, which can be converted to radio active elements by means of external irradiation after that anti-lymphoma antibody carrying said element has been accumulated to specific cells or tissues, or photoactive compounds or compounds used in photo imaging or photo dynamic therapy, or any other molecule having the same or similar effect, directly or indirectly, on lymphoma cells or lymphoma tissues.

11. The medical agent according to claim 10, wherein the effector agent comprises aryl halides and vinyl halides for radionuclides of halogens, N_2S_2 and N_3S chelates for Tc and Re radionuclides, amino-carboxy derivatives, preferably EDTA and DTPA or derivatives thereof, and cyclic amines, preferably NOTA, DOTA and TETA, and derivatives thereof, for In, Y, Pb, Bi, Cu, Sm and Lu radionuclides, or any other radionuclide capable of forming a complex with said chelates.

12. The medical agent according to claim 11, wherein the DTPA derivatives are Me-DTPA, CITC-DTPA, and cyclohexyl-DTPA.

13. The medical agent according to any one of claims 10-12, wherein the effector agent comprises DOTA and is provided with Y-90 for therapeutic application or In-111 for diagnostic application.

14. The medical agent according to any one of claims 10-13, wherein the effector agent is provided with positron imaging radionuclides, preferably F-18, Br-75, Br-76, and I-124; therapeutic radionuclides, preferably Y-90, I-131, In-114m, Re-186, Re-188, Cu-67, Sm-157, Lu-177, Bi-212, Bi-213, At-211, Ra-223; gamma imaging radionuclides, preferably Tc-99m, In-111, I-123, and I-125; beta radiation emitters, preferably scandium-46, scandium-47, scandium-48, copper-67, gallium-72, gallium-73, yttrium-90, ruthenium-97, palladium-100, rhodium-101,

palladium-109, samarium-153, lutetium-177, rhenium-186, rhenium-188, rhenium-189, gold-198, radium-212 and lead-212, gamma emitters, preferably iodine-131 and indium-114; and positron emitters, preferably gallium-68 and zirconium-89.

15. The medical agent according to claim 1, wherein the affinity ligand is capable of binding to another molecule having affinity for said ligand with an affinity binding constant of at least 10^6M^{-1} , preferably at least 10^8M^{-1} .

16. The medical agent according to claim 15, wherein the affinity ligand is a moiety which binds specifically to avidin, streptavidin or any other derivatives, mutants or fragments of avidin or streptavidin having essentially the same binding function to this affinity ligand.

17. The medical agent according to claim 16, wherein the affinity ligand is biotin, or a biotin derivative having essentially the same binding function to avidin or streptavidin as biotin.

18. The medical agent according to claim 17, wherein the biotin derivative is chosen from the group consisting of norbiotin, homobiotin, oxybiotin, iminobiotin, desti-biotin, diaminobiotin, biotin sulfoxide, and biotin sulfone, or derivatives thereof having essentially the same binding function.

19. The medical agent according to any one of claims 15-18, wherein the stability against enzymatic cleavage, preferably against cleavage by biotinidase, of the biotinamide bond to release biotin has been improved by using biotin derivatives, preferably norbiotin or homobiotin.

20. The medical agent according to claim 1, wherein the trifunctional cross-linking moiety is chosen from the group consisting of triaminobenzene, tricarboxybenzene, dicarboxyaniline and diaminobenzoic acid.

21. The medical agent according to claim 1, wherein linker 1 serves as an attaching moiety and a spacer between the trifunctional cross-linking moiety and the affinity ligand, preferably a biotin moiety, such that
5 binding with avidin or streptavidin, or any other biotin binding species, is not diminished by steric hindrance.

22. The medical agent according to claim 21, wherein linker 1 contains hydrogen bonding atoms, preferably ethers or tioethers, or ionisable groups, preferably
10 carboxylate, sulfonates, or ammonium groups to aid in water solubilisation of the biotin moiety.

23. The medical agent according to claim 22, wherein stability against enzymatic cleavage, preferably against cleavage by biotinidase, of the biotin amide bond to
15 release biotin has been provided by introducing an alpha carboxylate or an N-methyl group in linker 1.

24. The medical agent according to claim 1, wherein linker 2 provides a spacer length of 1-25 atoms, preferably a length of 6-18 atoms.

20 25. The medical agent according to claim 24, wherein linker 2 contains hydrogen bonding atoms, preferably ethers or tioethers, or ionisable groups, to aid in water solubilisation.

26. The medical agent according to claim 1, wherein
25 linker 2 is excluded.


27. The medical agent according to claim 1, wherein linker 3 provides a spacer of a length of 1-25 atoms, preferably a length of 6-18 atoms.

28. The medical agent according to claim 27, wherein
30 linker 3 contains hydrogen bonding atoms such as ethers or tioethers, or ionisable groups, preferably carboxylate, sulfonates, or ammonium groups to aid in water solubilisation.

29. The medical agent according to claim 1, wherein
35 linker 3 is excluded.

30. The medical agent according to any one of the preceding claims, wherein more than one affinity ligand and/or more than one effector agent are bound to a trifunctional or tetrafunctional cross-linking moiety.

31. The medical agent according to any one of claims 1-25 and 27-28, wherein it is



bound to the effector agent,

20 wherein the anti-CD20 antibody preferably is rituximab, wherein n is 2-4, preferably 3; o is 1-6, preferably 3; p is 1-6, preferably 3; R_2 is $=CH_2OH$ or $-CO_2H$; and R_1 is $-CH_3$, $-CH_2OH$, or $-H$.

25 32. The medical agent according to any one of the preceding claims, wherein it is 3-(13'-thiourea-benzyl-(DOTA)trioxadiazine-1-(13''-biotin-Asp-OH)trioxamine-5-isothio-cyanato-aminoisophthalate-ibritomumab, 3-(13'-thiourea-benzyl (DOTA)trioxadiazine-1-(13''-biotin-Asp-OH)trioxamine-5-isothio-cyanato-aminoisophthalate-rituximab, or 1-Isocyanato-3-((1S'-(N-Biotinyl)- β -L-Aspartyl)-4',7',10'-Trioxa-penta-Decanylamino)-1-((13-(Benzylthiourea-CHX-A'')-4,7,10-Trioxatridecanediamine)-Aminosio-phthalate-rituximab, preferably 3-(13'-thiourea-benzyl (DOTA)trioxadiazine-1-(13''-biotin-Asp-OH)trioxamine-5-isothio-cyanato-aminoisophthalate-rituximab.

33. A composition containing a medical agent according to any one of the preceding claims, wherein it further comprises physiologically acceptable additives, preferably an ammonium acetate solution.

5 34. A kit for extracorporeal elimination or at least reduction of the concentration of a non-tissue-bound therapeutic or diagnostic medical agent as defined in any one of claims 1-32 in the plasma or whole blood of a mammalian host, wherein said medical agent previously has
10 been introduced into a mammalian host and kept therein for a certain time in order to be concentrated to the specific tissues or cells by being attached thereto, said kit comprising

- a) the medical agent, and
- 15 b) an extracorporeal device comprising an immobilised receptor to which the affinity ligand adheres.

35. Use of a medical agent according to any one of claims 1-32 for the production of a medicament for the treatment of lymphoma, preferably non-Hodgkin's lymphoma.

20 36. A method for treatment of lymphoma, preferably non-Hodgkin's lymphoma, wherein anti-lymphoma antibodies or variants thereof as defined in any one of claims 2-6 are administered to a patient with a need thereof, wherein complexes formed between said anti-lymphoma antibodies
25 or variants thereof and leukocytes having said one or more cell surface antigen(s) are then eliminated from the body of the patient, followed by administration of the medical agent according to any one of claims 1-32, optionally together with said anti-lymphoma antibodies or
30 variants thereof as such, followed by extracorporeal elimination of the medical agent which has not been bound to the cell surface antigens on the lymphoma tumour cells.

37. The method according to claim 36, wherein the
35 effector agent of the medical agent is ^{90}Y and the medical agent is administered in a single dose of more than 20 MBq/kg body weight.

38. The method for diagnosing lymphoma, preferably non-Hodgkin's lymphoma, wherein anti-lymphoma antibodies or variants thereof as defined in any one of claims 2-6 are administered to a patient with a need thereof, where-
5 in complexes formed between said anti-lymphoma antibodies or variants thereof and leukocytes having said one or more cell surface antigen(s) are then eliminated from the body of the patient, followed by administration of the medical agent according to any one of claims 1-32, op-
10 tionally together with said anti-lymphoma antibodies or variants thereof as such, followed by extracorporeal elimination of the medical agent which has not been bound to the cell surface antigens on the lymphoma tumour cells.

39. The method according to claim 38, wherein the
15 effector agent of the medical agent is ^{90}Y or ^{111}In and the medical agent is administered in a dose range of 10-20, preferably 11-15, MBq/kg body weight in view of ^{90}Y and in a dose range of 20-250, preferably 50-150, MBq/kg body weight in view of ^{111}In .

20 40. A method for combined diagnosing and treatment of lymphoma, preferably non-Hodgkin's lymphoma, wherein a medical agent according to any one of claims 1-32, wherein the effector agent is ^{111}In , is administered to a patient in a dose range of 50-150 MBq/kg body weight, and a
25 medical agent according to any one of claims 1-32, wherein the effector agent is ^{90}Y , is administered to the patient in a dose of more than 20 MBq/kg body weight, either in sequence at intervals of 6-8 days or simultaneously.




Figure 1: Depletion of 1033-rituximab conjugates during re-circulation through a miniaturised Mitradep[®]. The lines display the theoretically maximal possible depletion rate. As seen the depletion of 1033-rituximab is not different from the theoretical depletion line, i.e. all 1033-rituximab present in the solution passing through the device is removed.

THIS PAGE BLANK (USPTO)

2/6




Figure 2: Flow cytometric assay of binding to the CD20-positive cell line

THIS PAGE BLANK (USPTO)

3/6




Figure 3: Binding of 1033-conjugates to a CD20+ (SB) and a CD20- (HSB) cell line. The binding is detected with an enzyme-labelled streptavidin.




Figure 4: Competitive inhibition of ^{125}I -labelled rituximab binding to SB cells by cold rituximab and 1033-rituximab conjugates.

JCO9 REC'd PCT/PTO 10 JUN 2001

THIS PAGE BLANK (USPTO)

4/6

4,6-1033-rituximab




Figure 5: Whole body clearance of radioactivity in rats injected with ^{111}In -1033-rituximab antibody conjugates expressed as percentage \pm std.dev. The data are corrected for radioactivity decay and background.

THIS PAGE BLANK (USPTO)

5/6




Figure 6: Blood clearance of ^{111}In -1033-rituximab antibody conjugates expressed as % injected dose/gram \pm std.dev. The data are corrected for radioactivity decay.




Figure 7: Biodistribution of ^{111}In -1033-rituximab (4.6 1033/IgG) in rats. The biodistribution is expressed as per cent of injected dose per gram tissue \pm std.dev. The results are corrected for radiochemical decay.

THIS PAGE BLANK (USPTO)

6/6




Figure 8: HPLC size exclusion separation of blood samples drawn from a rat injected with ^{111}In -1033-rituximab (4.6 1033/IgG).

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2003/001949

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: A61K 39/395, A61K 47/48, A61K 51/10 // A61M 1/36
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,OK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, BIOSIS, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 0002050 A1 (DEPARTMENT OF RADIATION ONCOLOGY, UNIVERSITY OF WASHINGTON), 13 January 2000 (13.01.2000), see pages 6-12, examples and claims, page 6, lines 16-36 --	1-10
Y	US 2001023288 A1 (WILBUR ET AL), 20 Sept 2001 (20.09.2001), see the claims --	1-40
A	Bioconjugate Chem., Volume 11, 2000, D. Scott Wibur et al, "Biotin Reagents for Antibody Pretargeting. 4. Selection of Biotin Conjugates for in Vivo Application Based on Their Dissociation Rate from Avidin and Streptavidin", pages 569-583 --	1-40

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

26 February 2004

Date of mailing of the international search report

03-03-2004

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

CARL-OLOF GUSTAFSSON/BS

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2003/001949

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Cancer Biotherapyh & Radiopharmaceuticals, Volume 15, No. 1, 2000, Paul L. Weiden et al, "Pretargeted Radioimmunotherapy (PRIT TM) for Treatment of Non-Hodgkin's Lymphoma (NHL): Initial Phase I/II Study Results", pages 15-29 --	1-40
Y	WO 0009160 A1 (IDEC PHARMACEUTICALS CORPORATION), 24 February 2000 (24.02.2000); page 12, line 24 - page 13, line 8; page 33, line 17 - line 34 --	1-40
Y	WO 0180884 A1 (IDEC PHARMACEUTICALS CORPORATION), 1 November 2001 (01.11.2001), page 37, line 29 - page 40 -- -----	1-40

INTERNATIONAL SEARCH REPORT

Information on patent family members

24/12/2003

International application No.

PCT/SE 2003/001949

WO	0002050	A1	13/01/2000	AU	5076799	A	24/01/2000
				AU	8366398	A	24/01/2000
				CA	2336739	A	13/01/2000
				EP	1095274	A	02/05/2001
				JP	2002519440	T	02/07/2002
				NO	20010021	A	07/03/2001
				SE	9801345	D	00/00/0000
				WO	0002051	A	13/01/2000

US	2001023288	A1	20/09/2001	AU	4161700	A	14/11/2000
				AU	5076799	A	24/01/2000
				CA	2336739	A	13/01/2000
				EP	1095274	A	02/05/2001
				EP	1173662	A	23/01/2002
				JP	2002519440	T	02/07/2002
				NO	20010021	A	07/03/2001
				SE	514444	C	26/02/2001
				SE	9901241	A	09/10/2000
				US	6655327	B	02/12/2003
WO	0061929	A	19/10/2000				

WO	0009160	A1	24/02/2000	AU	767965	B	27/11/2003
				AU	5553199	A	06/03/2000
				BR	9913645	A	25/09/2001
				CA	2340091	A	24/02/2000
				CN	1320044	T	31/10/2001
				CZ	20010526	A	13/11/2002
				EP	0865367	A	23/09/1998
				EP	1112084	A	04/07/2001
				HU	0103484	A	28/01/2002
				IL	141349	D	00/00/0000
				JP	2002522511	T	23/07/2002
				NO	20010699	A	10/04/2001
				PL	346046	A	14/01/2002
				US	6455043	B	24/09/2002
				US	2003026804	A	06/02/2003
				ZA	200101157	A	10/07/2002

WO	0180884	A1	01/11/2001	AU	5914201	A	07/11/2001
				CA	2405632	A	01/11/2001
				CN	1437478	T	20/08/2003
				EP	1286692	A	05/03/2003
				JP	2003531178	T	21/10/2003
				US	2002009444	A	24/01/2002

INTERNATIONAL SEARCH REPORT

International application No.
PCT SE 2003/001949

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **36-40**
because they relate to subject matter not required to be searched by this Authority, namely:
see extra sheet
2. ☒ Claims Nos.: **1**
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
see extra sheet
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Box II.1

Claims 36-40 relate to methods of treatment of the human or animal body by surgery or by therapy/diagnostic methods practised on the human or animal body/Rule 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.

The wording of claim 1 is ambiguous (see "1 d)") as the "biomolecule reactive moiety ... being capable of forming a bond with the ... antibody" is a reactive moiety that may be linked to the lymphoma antibody, while the claim pertains to an agent that comprises a reagent that is indeed conjugated to a lymphoma antibody. The search has been limited to agents with bound antilymphoma antibodies.

Present claims relate to a product/compound defined by reference to a desirable characteristic or property, namely comprising an affinity ligand, an effector agent and "a biomolecule reactive moiety". The claims cover all products/compounds having these characteristics or properties, whereas the application /provides support within the meaning of Article 6 PCT and / or disclosure within the meaning of Article 5 PCT for only a very limited number of such products/compounds. Additionally, previously known compounds may be included in the scope of the present claims. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product/compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the products/compounds identified in the examples and to the general aspects of the invention to the extent it was possible.

THIS PAGE BLANK (USPTO)